

A representation formula of the viscosity solution of the contact Hamilton-Jacobi equation and its applications*

Panrui Ni¹ Lin Wang² Jun Yan³

Abstract Assume M is a closed, connected and smooth Riemannian manifold. We consider the evolutionary Hamilton-Jacobi equation

$$\begin{cases} \partial_t u(x, t) + H(x, u(x, t), \partial_x u(x, t)) = 0, & (x, t) \in M \times (0, +\infty), \\ u(x, 0) = \varphi(x), \end{cases}$$

where $\varphi \in C(M)$ and the stationary one

$$H(x, u(x), \partial_x u(x)) = 0,$$

where $H(x, u, p)$ is continuous, convex and coercive in p , uniformly Lipschitz in u . By introducing a solution semigroup, we provide a *representation formula* of the viscosity solution of the evolutionary equation. As its applications, we obtain a necessary and sufficient condition for the existence of the viscosity solutions of the stationary equations. Moreover, we prove a new comparison theorem depending on the neighborhood of the projected Aubry set essentially, which is different from the one for the Hamilton-Jacobi equation independent of u .

Keywords Weak KAM theory, Hamilton-Jacobi equations, Aubry sets
2000 MR Subject Classification 37J50, 35F21, 35D40

1 Introduction and main results

The study of the theory of viscosity solutions of the following two forms of Hamilton-Jacobi equations

$$\partial_t u(x, t) + H(x, u(x, t), \partial_x u(x, t)) = 0, \quad (1.1)$$

and

$$H(x, u(x), \partial_x u(x)) = 0 \quad (1.2)$$

has a long history. There are many celebrated results on the existence, uniqueness, stability and large time behavior problems for the viscosity solutions (see [1, 9–11] for instance).

Manuscript received

¹Shanghai Center for Mathematical Sciences, Fudan University, Shanghai 200438, China.
E-mail: prni18@fudan.edu.cn

²School of Mathematics and Statistics, Beijing Institute of Technology, Beijing 100081, China.
E-mail: lwang@bit.edu.cn

³School of Mathematical Sciences, Fudan University, Shanghai 200433, China.
E-mail: yanjun@fudan.edu.cn

*Lin Wang is supported by NSFC Grant No. 12122109, 11790273. Jun Yan is supported by NSFC Grant No. 11790273, 11631006.

For the case with the Hamiltonian independent of u , its characteristic equation is the Hamilton equation. For the case with the Hamiltonian depending on u , the characteristic equation is called the contact Hamilton equation. In [23], the authors introduced an implicit variational principle for the contact Hamilton equation. Based on that, a representation formula was provided for the viscosity solution of the evolutionary equation in [24]. The existence of the solutions for the ergodic problem was also proved. In [25], the Aubry-Mather theory was developed for contact Hamiltonian systems with strictly increasing dependence on u . In [26], the authors further studied the strictly decreasing case, and discussed large time behavior of the viscosity solution of the evolutionary case.

In order to get the C^1 -regularity of the minimizers, it was assumed that $H(x, u, p)$ is of class C^3 in [23]. The results in [24–26] are based on the implicit variational principle established in [23]. Thus, all of them require the contact Hamiltonian to be of class C^3 . This paper is devoted to reducing the dynamical assumptions on the Hamiltonian: C^3 , *strictly convex* and *superlinear* to the standard PDE assumptions: *continuous*, *convex* and *coercive*. In this general case, the contact Hamiltonian equation can not be defined.

For the classical Hamilton-Jacobi (HJ) equation with time-independence, the related problems were considered in [12, 14]. Different from the previous works [12, 14], one has to face certain new difficulties due to the appearance of the Lavrentiev phenomenon caused by time-dependence. In addition, different from [24–26], we have not compactness of minimizers, which is a crucial ingredient in [24–26].

By combining dynamical and PDE approaches, we provide a *representation formula* of the viscosity solution of the evolutionary equation, which can be referred to as an implicit Lax-Oleinik semigroup. As its applications, we obtain a necessary and sufficient condition for the existence of the viscosity solutions of the stationary equations. It is well known that the comparison theorem plays a central role in the viscosity solution theory. We prove a new comparison result depending on a neighborhood of the projected Aubry set essentially. An example is constructed to show that the requirement of the neighborhood is *necessary* for a special class of Hamilton-Jacobi equations that do not satisfy the “proper” condition introduced in [9]. Comparably, the viscosity solution is determined completely by the projected Aubry set itself for the “proper” cases ([27, Theorem 1.6]).

Throughout this paper, we assume M is a closed (compact without boundary), connected and smooth Riemannian manifold and $H : T^*M \times \mathbb{R} \rightarrow \mathbb{R}$ satisfies

(C): $H(x, u, p)$ is continuous;

(CON): $H(x, u, p)$ is convex in p , for any $(x, u) \in M \times \mathbb{R}$;

(CER): $H(x, u, p)$ is coercive in p , i.e. $\lim_{\|p\| \rightarrow +\infty} (\inf_{x \in M} H(x, 0, p)) = +\infty$;

(LIP): $H(x, u, p)$ is Lipschitz in u , uniformly with respect to (x, p) , i.e., there exists $\lambda > 0$ such that $|H(x, u, p) - H(x, v, p)| \leq \lambda|u - v|$, for all $(x, p) \in T^*M$ and all $u, v \in \mathbb{R}$.

Correspondingly, one has the Lagrangian associated to H :

$$L(x, u, \dot{x}) := \sup_{p \in T_x^*M} \{\langle \dot{x}, p \rangle - H(x, u, p)\}.$$

A list of notations is provided at the end of this section.

Remark 1.1. Due to the absence of superlinearity of H , the Lagrangian L may take the value $+\infty$. Define

$$\text{dom}(L) := \{(x, \dot{x}, u) \in TM \times \mathbb{R} \mid L(x, u, \dot{x}) < +\infty\}.$$

Then L satisfies the following properties (see [12, Proposition 2.7] for instance)

- (LSC): $L(x, u, \dot{x})$ is lower semicontinuous, and continuous on the interior of $\text{dom}(L)$;
(CON): $L(x, u, \dot{x})$ is convex in \dot{x} , for any $(x, u) \in M \times \mathbb{R}$;
(LIP): $L(x, u, \dot{x})$ is Lipschitz in u , uniformly with respect to (x, \dot{x}) , i.e., there exists $\lambda > 0$ such that $|L(x, u, \dot{x}) - L(x, v, \dot{x})| \leq \lambda|u - v|$, for all $(x, \dot{x}, u) \in \text{dom}(L)$.

Remark 1.2.

- (1) The assumption (CER) is equivalent to the following statement: for each $R > 0$, there exists $K > 0$ such that for any $|u| < R$ and $\|p\| > K$, we have $H(x, u, p) > R$. In fact, by (CER), for each $R > 0$, there exists $K > 0$ such that for $\|p\| > K$, $H(x, 0, p) > (1 + \lambda)R$. By (LIP), for any $|u| < R$,

$$H(x, u, p) \geq H(x, 0, p) - \lambda|u| > R.$$

The converse direction is obvious.

- (2) $\text{dom}(L)$ is independent of u . More precisely, given $(x, \dot{x}) \in TM$, if $L(x, u_0, \dot{x}) < +\infty$ for a given $u_0 \in \mathbb{R}$, then for any $u \in \mathbb{R}$,

$$\begin{aligned} L(x, u, \dot{x}) &\leq \sup_{p \in T_x^* M} \{\langle \dot{x}, p \rangle - H(x, u_0, p)\} + \lambda|u - u_0| \\ &= L(x, u_0, \dot{x}) + \lambda|u - u_0| < +\infty. \end{aligned}$$

Thus, there holds

$$\text{dom}(L) = \{(x, \dot{x}) \in TM \mid L(x, 0, \dot{x}) < +\infty\} \times \mathbb{R}.$$

1.1 An implicit Lax-Oleinik semigroup

Consider the viscosity solution of the Cauchy problem

$$\begin{cases} \partial_t u(x, t) + H(x, u(x, t), \partial_x u(x, t)) = 0, & (x, t) \in M \times (0, +\infty). \\ u(x, 0) = \varphi(x), & x \in M. \end{cases} \quad (CP_H)$$

We have the following result.

Theorem 1. Assume $H : T^*M \times \mathbb{R} \rightarrow \mathbb{R}$ satisfies (C)(CON)(CER)(LIP). The following implicit backward Lax-Oleinik semigroup $T_t^- : C(M) \rightarrow C(M)$, via

$$T_t^- \varphi(x) = \inf_{\gamma(\dot{t})=x} \left\{ \varphi(\gamma(0)) + \int_0^t L(\gamma(\tau), T_\tau^- \varphi(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau \right\} \quad (T^-)$$

is well-defined. The infimum is taken among absolutely continuous curves $\gamma : [0, t] \rightarrow M$ with $\gamma(t) = x$. Moreover,

- (i) if the initial condition φ is continuous, then $u(x, t) := T_t^- \varphi(x)$ represents the unique continuous viscosity solution of (CP_H) .
- (ii) If φ is Lipschitz continuous, then $u(x, t) := T_t^- \varphi(x)$ is also locally Lipschitz continuous on $M \times [0, +\infty)$.

The main difficulties to prove Theorem 1 are stated as follows.

- Compared to the contact HJ equation under the Tonelli conditions, the contact Hamilton flow can not be defined. Consequently, we do not have the compactness of the minimizing orbit set, which plays a crucial role in the previous work on contact HJ equations (see [24, Lemma 2.1]).
- Compared to the classical HJ equation in less regular cases (see [12, 14]), the backward Lax-Oleinik semigroup is implicit defined, which causes t -dependence of the Lagrangian. Due to the Lavrentiev phenomenon, it is not direct to prove the Lipschitz continuity of the minimizers of $T_t^- \varphi(x)$ (see [2] for various counterexamples).

Under the general assumptions (C) (CON) (CER) and (LIP), the *Lipschitz continuity* of $T_t^- \varphi(x)$ and its minimizers are proved by combining dynamical and PDE approaches, together with a new variational inequality introduced in [3].

Remark 1.3. *Similar to Theorem 1, the forward Lax-Oleinik semigroup can be defined as*

$$T_t^+ \varphi(x) = \sup_{\gamma(0)=x} \left\{ \varphi(\gamma(t)) - \int_0^t L(\gamma(\tau), T_{t-\tau}^+ \varphi(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau \right\}. \quad (\text{T+})$$

Use the same argument as [25, Proposition 2.8], one has $T_t^+ \varphi := -\bar{T}_t^-(-\varphi)$, where \bar{T}_t^- denotes the backward Lax-Oleinik semigroup associated to $L(x, -u, -\dot{x})$.

By Theorem 1, if the fixed points of T_t^- exist, then they are viscosity solutions of

$$H(x, u(x), \partial_x u(x)) = 0. \quad (E_H)$$

Recently, an alternative variational formulation was provided in [6, 7, 21] in light of G. Herglotz's work [15], which is related to nonholonomic constraints. By using the Herglotz variational principle, various kinds of representation formulae for the viscosity solutions of (1.1) were also obtained in [16]. For simplicity, we will omit the word ‘‘viscosity’’ if it is not necessary to be mentioned.

1.2 An existence result for the solutions of (E_H)

Remark 1.4. *Let $H : T^*M \times \mathbb{R} \rightarrow \mathbb{R}$ satisfy (C)(CER)(LIP). According to the Perron method [17], if (E_H) has a Lipschitz subsolution f and a Lipschitz supersolution g such that $f \leq g$. Then the equation (E_H) admits a Lipschitz viscosity solution.*

In light of [17], we introduce another necessary and sufficient condition for (E_H) to admit solutions.

Theorem 2. *Let $H : T^*M \times \mathbb{R} \rightarrow \mathbb{R}$ satisfy (C)(CON)(CER)(LIP). The following statements are equivalent:*

- (1) (E_H) admits Lipschitz solutions;
- (2) There exist two continuous functions φ and ψ such that $T_t^- \varphi \geq C_1$ and $T_t^- \psi \leq C_2$, where C_1, C_2 are constant independent of t and x ;
- (3) There exist two continuous functions φ and ψ , and $t_1, t_2 > 0$ such that $T_{t_1}^- \varphi \geq \varphi$ and $T_{t_2}^- \psi \leq \psi$.

If (E_H) admits a solution u , one can take u as the initial function. The statement (2) and (3) hold true obviously. Thus, we only need to show the opposite direction, which will be proved in Section 3. The main novelty of Theorem 2 is that the lower bound of $T_t^- \varphi$ is not required to be less than or equal to the upper bound of $T_t^- \psi$.

1.3 The Aubry set

We denote by \mathcal{S}_- and \mathcal{S}_+ the set of all backward weak KAM solutions and the set of all forward weak KAM solutions of (E_H) respectively. See Appendix D for their definitions and relations with viscosity solutions. In the discussion below, we need to introduce the following assumption

(S): The set \mathcal{S}_- is nonempty. Namely, (E_H) admits a solution.

Definition 1.5. Let $u_- \in \mathcal{S}_-$, $u_+ \in \mathcal{S}_+$. We define the projected Aubry set with respect to u_- by

$$\mathcal{I}_{u_-} := \{x \in M : u_-(x) = \lim_{t \rightarrow +\infty} T_t^+ u_-(x)\}.$$

Similarly, we define the projected Aubry set with respect to u_+ by

$$\mathcal{I}_{u_+} := \{x \in M : u_+(x) = \lim_{t \rightarrow +\infty} T_t^- u_+(x)\}.$$

In particular, if $u_+(x) = \lim_{t \rightarrow +\infty} T_t^+ u_-(x)$ and $u_-(x) = \lim_{t \rightarrow +\infty} T_t^- u_+(x)$, then

$$\mathcal{I}_{u_-} = \mathcal{I}_{u_+},$$

which is denoted by $\mathcal{I}_{(u_-, u_+)}$, following the notation introduced by Fathi [13].

Theorem 3. Assume $H : T^*M \times \mathbb{R} \rightarrow \mathbb{R}$ satisfies (C)(CON)(CER)(LIP) and (S). Let $u_- \in \mathcal{S}_-$. Then

- (1) the limit function $\lim_{t \rightarrow +\infty} T_t^+ u_-(x)$ exists and it is a forward weak KAM solution. For each $u_+ \in \mathcal{S}_+$, the limit function $\lim_{t \rightarrow +\infty} T_t^- u_+(x)$ exists and it is a backward weak KAM solution of (E_H) ;
- (2) both \mathcal{I}_{u_-} and \mathcal{I}_{u_+} are nonempty.

By Remark 1.3, we only need to prove Theorem 3 for $T_t^+ u_-(x)$ and \mathcal{I}_{u_-} .

1.4 A comparison result for the solutions of (E_H)

In this part, we are concerned with further properties of the viscosity solution for a special class of Hamilton-Jacobi equations that do not satisfy the proper condition:

$$H(x, r, p) \leq H(x, s, p) \quad \text{whenever } r \leq s.$$

We assume $H : T^*M \times \mathbb{R} \rightarrow \mathbb{R}$ satisfies (C), (CON), (CER), (LIP) and

(STD): $H(x, u, p)$ is strictly decreasing in u .

Under the assumptions above, the solution of $H(x, u, \partial_x u) = 0$ is not unique (see e.g., Example (E1) below). The following result provides a comparison among different solutions.

Theorem 4. *Let $v_1, v_2 \in \mathcal{S}_-$.*

- (1) *If $v_1 \leq v_2$, then $\emptyset \neq \mathcal{I}_{v_2} \subseteq \mathcal{I}_{v_1}$;*
- (2) *If there is a neighborhood \mathcal{O} of \mathcal{I}_{v_2} such that $v_1|_{\mathcal{O}} \leq v_2|_{\mathcal{O}}$, then $v_1 \leq v_2$ everywhere;*
- (3) *If $\mathcal{I}_{v_1} = \mathcal{I}_{v_2}$ and $v_1|_{\mathcal{O}} = v_2|_{\mathcal{O}}$, then $v_1 = v_2$ everywhere.*

In order to explain the necessity of the neighbourhood \mathcal{O} , we consider the following example

$$-\lambda u(x) + \frac{1}{2}|u'(x)|^2 + V(x) = 0, \quad x \in \mathbb{S} \simeq (-1, 1], \quad (\text{E1})$$

where \mathbb{S} denotes a flat circle with the fundamental domain $(-1, 1]$, and $V(x)$ is the restriction of $x^2/2$ on \mathbb{S} . Then $H(x, u, p) = -\lambda u + |p|^2/2 + V(x)$ defined on $T^*\mathbb{S} \times \mathbb{R}$ is Lipschitz continuous. Assume $\lambda > 2$, then two viscosity solutions of (E1) are

$$u_1(x) = \frac{\lambda + \sqrt{\lambda^2 - 4}}{2}V(x), \quad u_2(x) = \frac{\lambda - \sqrt{\lambda^2 - 4}}{2}V(x).$$

It can be shown that $\mathcal{I}_{u_1} = \mathcal{I}_{u_2} = \{0\}$, although $u_1 \neq u_2$ on \mathbb{S} . A detailed analysis of Example (E1) is given in Section 6 below.

The rest of this paper is organized as follows. In Section 2, we prove Theorem 1. To achieve that, we need some technical lemmas whose proofs are given in Appendix B and C. Theorem 2, Theorem 3 and Theorem 4 are proved in Section 3, Section 4 and Section 5 successively. In addition, we give some basic results on the existence and regularity of the minimizers of one dimensional variational problems in Appendix A, and we also provide some basic properties of weak KAM solution and viscosity solution in Appendix D for the reader's convenience.

We list notations in the present paper:

- $\text{diam}(M)$ denotes the diameter of M ;
- $d(x, y)$ denotes the distance between x and y induced by the Riemannian metric g on M ;
- $\|\cdot\|$ denotes the norms induced by g on both tangent and cotangent spaces of M ;
- $B(v, r)$ stands for the open norm ball on $T_x M$ centered at $v \in T_x M$ with radius r , and $\bar{B}(v, r)$ stands for its closure;
- $C(M)$ stands for the space of continuous functions on M ;
- $\text{Lip}(M)$ stands for the space of Lipschitz continuous functions on M ;
- $\|\cdot\|_\infty$ stands for the supremum norm of the vector-valued function on its domain.

2 An implicit Lax-Oleinik semigroup

In this part, we are devoted to proving Theorem 1. It is needed to show

- (*) if the initial condition φ is Lipschitz continuous, then $u(x, t) := T_t^- \varphi(x)$ is the Lipschitz solution of (CP_H) ;
- (**) if φ is continuous, then $u(x, t) := T_t^- \varphi(x)$ is the continuous solution of (CP_H) .

2.1 On Item (*): the Lipschitz initial condition

For the reader's convenience, we give a sketch of proof of Item (*) as follows.

- (1) Lemma 2.1 is proved in Appendix B. Some background knowledge is given in Section A.1.
- (2) Under **Condition 1**:
 u_k defined in (2.1) is continuous on $M \times [0, T]$ for each $k \in \mathbb{N}_+$,
 we prove Lemma 2.2(i) based on Lemma 2.1.
- (3) Under **Condition 2**:
 u_k is locally Lipschitz on $M \times (0, T]$ for each $k \in \mathbb{N}_+$, and it is the solution of (2.2) below,
 we prove Lemma 2.2(ii) by Lemma 2.2(i).
- (4) After a *superlinear* modification, we have Lemma 2.3 whose proof is provided in Appendix C. The proof needs some ingredients given in Section A.2.
- (5) **Condition 2** in Lemma 2.2(ii) can be verified by Lemma 2.3 under the *superlinear* condition. By Lemma 2.2(ii), Item (*) holds under the *superlinear* condition.
- (6) Under the *coercive* condition, we prove Lemma 2.4 based on Item (*) under the *superlinear* condition.
- (7) Under the *coercive* condition, **Condition 2** in Lemma 2.2(ii) can be verified by Lemma 2.4. By Lemma 2.2(ii), Item (*) holds under the *coercive* condition.

The following is a detailed proof of Item (*).

Lemma 2.1. Fix $T > 0$. Given $\varphi \in C(M, \mathbb{R})$, $v \in C(M \times [0, T], \mathbb{R})$ and $t \in [0, T]$, the functional

$$\mathbb{L}^t(\gamma) := \varphi(\gamma(0)) + \int_0^t L(\gamma(s), v(\gamma(s), s), \dot{\gamma}(s)) ds$$

reaches its infimum in the class of curves

$$X_t(x) = \{\gamma \in W^{1,1}([0, t], M) : \gamma(t) = x\}.$$

Lemma 2.2. Fix $T > 0$ and $\varphi \in C(M)$. For $k \in \mathbb{N}_+$ and $t \in (0, T]$, consider the following iteration procedure

$$u_k(x, t) := \inf_{\gamma(t)=x} \left\{ \varphi(\gamma(0)) + \int_0^t L(\gamma(\tau), u_{k-1}(\gamma(\tau), \tau), \dot{\gamma}(\tau)) d\tau \right\}, \quad (2.1)$$

where $u_0(x, t) := \varphi(x)$.

(i) If u_k is continuous on $M \times [0, T]$ for each $k \in \mathbb{N}_+$, then $\{u_k(x, t)\}_{k \in \mathbb{N}}$ converges uniformly to $u(x, t) := T_t^- \varphi(x)$ for all $(x, t) \in M \times [0, T]$, where the semigroup $T_t^- : C(M) \rightarrow C(M)$ is formulated as (T^-) .

(ii) Let $\varphi \in \text{Lip}(M)$. If u_k is locally Lipschitz on $M \times (0, T]$ for each $k \in \mathbb{N}_+$, and it is the solution of

$$\begin{cases} \partial_t u(x, t) + H(x, u_{k-1}(x, t), \partial_x u(x, t)) = 0, \\ u(x, 0) = \varphi(x), \end{cases} \quad (2.2)$$

then u_k is Lipschitz on $M \times [0, T]$, and its Lipschitz constant depends only on $\sup_{k \in \mathbb{N}} \|u_k\|_\infty$ and $\|\partial_x \varphi\|_\infty$. Moreover, the limit function $u(x, t) := T_t^- \varphi$ is Lipschitz.

Proof. Item (i): By Lemma 2.1, the minimizers of each u_k exist. First, let $\gamma_1 : [0, t] \rightarrow M$ be a minimizer of $u_1(x, t)$, then

$$\begin{aligned} u_2(x, t) - u_1(x, t) &\leq \int_0^t \left[L(\gamma_1(s), u_1(\gamma_1(s), s), \dot{\gamma}_1(s)) - L(\gamma_1(s), \varphi(\gamma_1(s)), \dot{\gamma}_1(s)) \right] ds \\ &\leq \lambda \|u_1 - \varphi\|_\infty t. \end{aligned}$$

Exchanging the positions of u_2 and u_1 , we obtain

$$|u_2(x, t) - u_1(x, t)| \leq \lambda \|u_1 - \varphi\|_\infty t.$$

Let $\gamma_2 : [0, t] \rightarrow M$ be a minimizer of $u_2(x, t)$, then

$$\begin{aligned} u_3(x, t) - u_2(x, t) &\leq \int_0^t \left[L(\gamma_2(s), u_2(\gamma_2(s), s), \dot{\gamma}_2(s)) - L(\gamma_2(s), u_1(\gamma_2(s), s), \dot{\gamma}_2(s)) \right] ds \\ &\leq \lambda \int_0^t |u_2(\gamma_2(s), s) - u_1(\gamma_2(s), s)| ds \leq \lambda \int_0^t \lambda \|u_1 - \varphi\|_\infty s ds \\ &= \frac{(\lambda t)^2}{2} \|u_1 - \varphi\|_\infty. \end{aligned}$$

Exchanging the positions of u_3 and u_2 , we obtain

$$|u_2(x, t) - u_1(x, t)| \leq \frac{(\lambda t)^2}{2} \|u_1 - \varphi\|_\infty.$$

Continuing the above procedure, we obtain

$$\|u_{j+1} - u_j\|_\infty \leq \frac{(\lambda T)^j}{j!} \|u_1 - \varphi\|_\infty,$$

for each $j \in \mathbb{N}$. Thus,

$$\|u_k - \varphi\|_\infty \leq \sum_{j=0}^{k-1} \|u_{j+1} - u_j\|_\infty \leq \sum_{j=0}^{k-1} \frac{(\lambda T)^j}{j!} \|u_1 - \varphi\|_\infty \leq e^{\lambda T} \|u_1 - \varphi\|_\infty, \quad \forall k \in \mathbb{N}_+.$$

For $k_1 > k_2$, we have

$$\|u_{k_1} - u_{k_2}\|_\infty \leq \frac{(\lambda T)^{k_2}}{k_2!} \|u_{k_1 - k_2} - \varphi\|_\infty \leq \frac{(\lambda T)^{k_2}}{k_2!} e^{\lambda T} \|u_1 - \varphi\|_\infty.$$

Since $(\lambda T)^k/k!$ converges to zero as $k \rightarrow \infty$, the right hand side can be arbitrarily small when k_2 is large enough. Therefore, the sequence $\{u_k(x, t)\}_{k \in \mathbb{N}}$ is a Cauchy sequence in the Banach space $(C(M \times [0, T]), \|\cdot\|_\infty)$. Then $\{u_k(x, t)\}_{k \in \mathbb{N}}$ converges uniformly to a continuous function $u(x, t)$. Define $\mathcal{A}_\varphi : C(M \times [0, T]) \rightarrow C(M \times [0, T])$ via

$$\mathcal{A}_\varphi[u](x, t) := \inf_{\gamma(t)=x} \left\{ \varphi(\gamma(0)) + \int_0^t L(\gamma(\tau), u(\gamma(\tau), \tau), \dot{\gamma}(\tau)) d\tau \right\}.$$

Then the limit function $u(x, t)$ satisfies

$$\|\mathcal{A}_\varphi[u] - u\|_\infty \leq \|\mathcal{A}_\varphi[u] - u_k\|_\infty + \|u_k - u\|_\infty \leq \lambda T \|u - u_{k-1}\|_\infty + \|u_k - u\|_\infty.$$

Setting $k \rightarrow +\infty$ we conclude that $u(x, t)$ is the unique fixed point of \mathcal{A}_φ . Namely, $u(x, t) := T_t^- \varphi$. The semigroup property of T_t^- can be verified by a similar argument as [20, Proposition 3.3].

Item (ii): By Item (i), $\{u_k(x, t)\}_{k \in \mathbb{N}}$ converges uniformly, then

$$\sup_{k \in \mathbb{N}} \|u_k(x, t)\|_\infty < +\infty.$$

Define

$$K_1 := \max\{|H(x, u, p)| : x \in M, |u| \leq \sup_{k \in \mathbb{N}} \|u_k(x, t)\|_\infty, \|p\| \leq \|\partial_x \varphi(x)\|_\infty\},$$

and

$$K_2 := \min\{H(x, u, p) : (x, p) \in T^*M, |u| \leq \sup_{k \in \mathbb{N}} \|u_k(x, t)\|_\infty\}.$$

We will prove by induction

$$\|\partial_t u_k(\cdot, t)\|_\infty \leq \max\{K_1 e^{\lambda t}, |K_2|\} \quad (2.3)$$

for each $k \in \mathbb{N}_+$.

First, let us consider the case $k = 1$. Define

$$K_0 := \max\{|H(x, u, p)| : x \in M, |u| \leq \|\varphi(x)\|_\infty, \|p\| \leq \|\partial_x \varphi(x)\|_\infty\}.$$

For any $h > 0$, we define

$$w(x, t) := \begin{cases} \varphi(x) - K_0 t, & t \leq h, \\ u_k(x, t - h) - K_0 h, & t > h. \end{cases} \quad (2.4)$$

First, the Lipschitz function $(x, t) \mapsto \varphi(x) - K_0 t$ satisfies

$$\partial_t u + H(x, \varphi(x), \partial_x u) \leq 0 \quad (2.5)$$

almost everywhere. According to Proposition D.3, $w(x, t)$ is a subsolution of (2.5) for $t \leq h$. Also, $w(x, t)$ satisfies (2.5) for $t > h$. Thus, $w(x, t)$ is a continuous subsolution of (2.5) with $w(x, 0) = \varphi(x)$. By the comparison result (see, e.g., [1, Theorem 5.1]), since $(x, t) \mapsto \varphi(x) - K_0 t$ is Lipschitz in x , we have

$$w(x, h) = \varphi(x) - K_0 h \leq u_1(x, h).$$

Note that $u_1(x, t)$ is Lipschitz on $M \times [h, T]$, we have

$$u_1(x, t) - K_0 h = w(x, t + h) \leq u_1(x, t + h), \quad \forall t \geq 0, h > 0.$$

Let $h \rightarrow 0^+$. We have $\partial_t u_1(x, t) \geq -K_0 \geq -K_1$. We also have

$$\partial_t u_1(x, t) = -H(x, \varphi(x), \partial_x u_1(x, t)) \leq |K_2|.$$

Thus, (2.3) holds for $k = 1$.

Now assume (2.3) holds for $k - 1$. For any $h > 0$, we define

$$\bar{w}(x, t) := \begin{cases} \varphi(x) - K_1 e^{\lambda h} t, & t \leq h, \\ u_k(x, t - h) - K_1 h e^{\lambda t}, & t > h. \end{cases} \quad (2.6)$$

First, the Lipschitz function $(x, t) \mapsto \varphi(x) - K_1 e^{\lambda h} t$ satisfies

$$\partial_t u + H(x, u_{k-1}(x, t), \partial_x u) \leq 0$$

almost everywhere. According to Proposition D.3, $\bar{w}(x, t)$ is a subsolution of (2.2) for $t \leq h$. For $t > h$, we have

$$\begin{aligned} & \partial_t \bar{w}(x, t) + H(x, u_{k-1}(x, t), \partial_x \bar{w}(x, t)) \\ &= \partial_t u_k(x, t - h) - K_1 h \lambda e^{\lambda t} + H(x, u_{k-1}(x, t), \partial_x u_k(x, t - h)) \\ &\leq \partial_t u_k(x, t - h) - \lambda \sup_{s \in [t-h, t]} \|\partial_t u_{k-1}(\cdot, s)\|_\infty h + H(x, u_{k-1}(x, t), \partial_x u_k(x, t - h)) \\ &\leq \partial_t u_k(x, t - h) + H(x, u_{k-1}(x, t - h), \partial_x u_k(x, t - h)) = 0. \end{aligned}$$

By the comparison result, since $(x, t) \mapsto \varphi(x) - K_1 e^{\lambda h} t$ is Lipschitz in x , we have

$$\bar{w}(x, h) = \varphi(x) - K_1 h e^{\lambda h} \leq u_k(x, h).$$

Note that $u_k(x, t)$ is Lipschitz on $M \times [h, T]$, we have

$$u_k(x, t) - K_1 h e^{\lambda(t+h)} = \bar{w}(x, t + h) \leq u_k(x, t + h), \quad \forall t \geq 0, h > 0.$$

Let $h \rightarrow 0^+$. We have $\partial_t u_k(x, t) \geq -K_1 e^{\lambda t}$. We also have

$$\partial_t u_k(x, t) = -H(x, u_{k-1}(x, t), \partial_x u_k(x, t)) \leq |K_2|.$$

We conclude that (2.3) holds for k . Plugging them into (2.2), one obtain

$$H(x, 0, \partial_x u_k(x, t)) \leq \max\{K_1 e^{\lambda T}, |K_2|\} + \lambda \|u_{k-1}(x, t)\|_\infty.$$

Thus $\|\partial_x u_k(x, t)\|_\infty$ is bounded on $M \times [0, T]$ by (CER). It means $u_k(x, t)$ is Lipschitz on $M \times [0, T]$, and the Lipschitz constant only depends on $\sup_{k \in \mathbb{N}} \|u_k(x, t)\|_\infty$ and $\|\partial_x \varphi(x)\|_\infty$. Moreover, $\{u_k(x, t)\}_{k \in \mathbb{N}}$ is equi-Lipschitz with respect to k . It follows that the limit function $u(x, t) := T_t^- \varphi$ is Lipschitz. \square

According to Lemma 2.2 (ii), the key point for the proof of Item (*) is to show for each $k \in \mathbb{N}$, $u_k(x, t)$ defined by (2.1) is locally Lipschitz, and solves (2.2) in the viscosity sense. We divide the remaining proof into two steps. In Step 1, we prove Item (*) for the Hamiltonian $H(x, u, p)$ depending on p superlinearly. In Step 2, the superlinearity is relaxed to (CER).

2.1.1 Step 1: Proof under the superlinear condition

In this part, we assume the Hamiltonian $H : T^*M \times \mathbb{R} \rightarrow \mathbb{R}$ satisfies (C)(CON)(LIP) and the following.

(SL): For every $(x, u) \in M \times \mathbb{R}$, $H(x, u, p)$ is superlinear in p , i.e. there exists a function $\Theta : [0 + \infty) \rightarrow \mathbb{R}$ satisfying

$$\lim_{r \rightarrow +\infty} \frac{\Theta(r)}{r} = +\infty, \quad \text{and} \quad H(x, u, p) \geq \Theta(\|p\|) \quad \text{for every } (x, p, u) \in T^*M \times \mathbb{R}.$$

The corresponding Lagrangian satisfies (CON)(LIP) and

(C): $L(x, u, \dot{x})$ is continuous;

(SL): For every $(x, u) \in M \times \mathbb{R}$, $L(x, u, \dot{x})$ is superlinear in \dot{x} , i.e., there exists a function $\Theta : [0 + \infty) \rightarrow \mathbb{R}$ satisfying

$$\lim_{r \rightarrow +\infty} \frac{\Theta(r)}{r} = +\infty, \quad \text{and} \quad L(x, u, \dot{x}) \geq \Theta(\|\dot{x}\|) \quad \text{for every } (x, \dot{x}, u) \in TM \times \mathbb{R}.$$

At the beginning, we need some technical results.

Lemma 2.3. *Given $T > 0$ and $\varphi \in C(M)$, if $v(x, t)$ is a Lipschitz function on $M \times [0, T]$, then*

(1) *for any $(x, t) \in M \times [0, T]$, the minimizers of*

$$u(x, t) := \inf_{\gamma(t)=x} \left\{ \varphi(\gamma(0)) + \int_0^t L(\gamma(\tau), v(\gamma(\tau), \tau), \dot{\gamma}(\tau)) d\tau \right\} \quad (2.7)$$

are Lipschitz. For any $r > 0$, if $d(x, x') \leq r$ and $|t - t'| \leq r/2$, where $t \geq r > 0$, then the Lipschitz constants of the minimizers of $u(x', t')$ only depend on (x, t) and r .

(2) *the value function $u(x, t)$ defined in (2.7) is locally Lipschitz on $M \times (0, T]$.*

(3) *$u(x, t)$ is also the viscosity solution of*

$$\begin{cases} \partial_t u(x, t) + H(x, v(x, t), \partial_x u(x, t)) = 0, \\ u(x, 0) = \varphi(x). \end{cases} \quad (2.8)$$

on $M \times [0, T]$.

Based on Lemma 2.3, we verify Item (*) under the assumption (SL). In fact, let $u_0 := \varphi \in \text{Lip}(M)$ in the iteration procedure given by (2.1). By Lemma 2.2 (i), $u_k(x, t)$ converges uniformly to $u(x, t) := T_t^- \varphi(x)$ on $M \times [0, T]$. By Lemma 2.3 (2) and (3), $u_1(x, t)$ satisfies the assumptions in Lemma 2.2 (ii), by which u_1 is Lipschitz on $M \times [0, T]$. Repeating the argument, one can obtain that u_k is the Lipschitz solution of (2.2). By Lemma 2.2 (ii), the Lipschitz constant of $u_k(x, t)$ is uniform with respect to k on $M \times [0, T]$. Since $H_k(t, x, p) := H(x, u_k(x, t), p)$ converges uniformly on compact subsets of $\mathbb{R} \times T^*M$, and $u_k(x, t)$ converges uniformly on $M \times [0, T]$, then the backward semigroup $u(x, t) := T_t^- \varphi(x)$, as the limit of $u_k(x, t)$, is the Lipschitz solution of (CP_H) by the stability of viscosity solutions.

2.1.2 Step 2: Relaxed to the coercive condition

In this part, we assume the Hamiltonian $H : T^*M \times \mathbb{R} \rightarrow \mathbb{R}$ satisfies (C)(CON)(CER)(LIP). By Lemma 2.1, one has the existence of the minimizers. In order to obtain the Lipschitz regularity of u_k in (2.1). We make a modification:

$$H_n(x, u, p) := H(x, u, p) + \max\{\|p\|^2 - n^2, 0\}, \quad n \in \mathbb{N}.$$

It is clear that H_n is superlinear in p . The sequence H_n is decreasing, and converges uniformly to H on compact subsets of $T^*M \times \mathbb{R}$. The sequence of the corresponding Lagrangians $\{L_n\}_{n \in \mathbb{N}}$ is increasing, and converges to L pointwisely. Denote by $u_{n,k}(x, t)$ the solution of (2.2) with H replaced by H_n .

Lemma 2.4. *Let H satisfy (C)(CON)(CER)(LIP). Let L be the Lagrangian associated to H . Given $\varphi \in \text{Lip}(M)$, for each $k \in \mathbb{N}$, the function $u_k(x, t)$ defined by (2.1) is the Lipschitz solution of (2.2).*

Proof. Given $n \in \mathbb{N}$, let

$$u_{n,k}(x, t) := \inf_{\gamma(t)=x} \left\{ \varphi(\gamma(0)) + \int_0^t L_n(\gamma(\tau), u_{n,k-1}(\gamma(\tau), \tau), \dot{\gamma}(\tau)) d\tau \right\}, \quad (2.9)$$

with $u_{n,0} := \varphi \in \text{Lip}(M)$. We first prove the following assertion for each $k \in \mathbb{N}$ by induction.

A[k]: Fix $k \in \mathbb{N}$. The sequence $\{u_{n,k}(x, t)\}_{n \in \mathbb{N}}$ is uniformly bounded and equi-Lipschitz with respect to n , and converges uniformly to $u_k(x, t)$ on $M \times [0, T]$. Moreover, the limit function $u_k(x, t)$ is Lipschitz.

We first prove that the assertion A[1] holds. By definition of H_n , for $n \geq \|\partial_x \varphi(x)\|_\infty$,

$$K_0^n := \max\{|H_n(x, u, p)| : x \in M, |u| \leq \|\varphi(x)\|_\infty, \|p\| \leq \|\partial_x \varphi(x)\|_\infty\}$$

is always equal to

$$K_0 = \max\{|H(x, u, p)| : x \in M, |u| \leq \|\varphi(x)\|_\infty, \|p\| \leq \|\partial_x \varphi(x)\|_\infty\}.$$

Note that $u_{n,1}(x, t)$ is the solution of

$$\partial_t u + H_n(x, \varphi(x), \partial_x u) = 0. \quad (2.10)$$

Similar to the proof of (2.3) with $k = 1$, we have

$$\partial_t u_{n,1}(x, t) \geq -K_0^n.$$

Combining (2.10) and the definition of H_n , we have

$$H(x, 0, \partial_x u_{n,1}(x, t)) \leq H_n(x, 0, \partial_x u_{n,1}(x, t)) \leq K_0 + \lambda \|\varphi(x)\|_\infty, \quad \forall n \geq \|\partial_x \varphi(x)\|_\infty.$$

Therefore, $\{u_{n,1}(x, t)\}_{n \in \mathbb{N}}$ is equi-Lipschitz. Note that

$$u_{n,1}(x, 0) = \varphi(x).$$

It follows that $\{u_{n,1}(x, t)\}_{n \in \mathbb{N}}$ is uniformly bounded, so it has a converging subsequence. According to Lemma A.7, $u_{n,1}(x, t)$ converges to $u_1(x, t)$ pointwisely. It follows that

$$\lim_{n \rightarrow +\infty} u_{n,1}(x, t) = u_1(x, t), \quad \text{uniformly on } M \times [0, T],$$

which implies that $u_1(x, t)$ is Lipschitz.

Now assume that the assertion A[k-1] holds. Then $u_{k-1}(x, t)$ is Lipschitz, and $l_{k-1} := \sup_{n \in \mathbb{N}} \|u_{n,k-1}(x, t)\|_\infty$ is finite. We will prove A[k] from A[k-1]. First, plugging the continuous function $u_{k-1}(x, t)$ into (2.1) and by Lemma 2.1, the minimizers of $u_k(x, t)$ exist in the class of absolutely continuous curves. By definition of H_n , for $n \geq \|\partial_x \varphi(x)\|_\infty$,

$$K_n := \max\{|H_n(x, u, p)| : x \in M, |u| \leq l_{k-1}, \|p\| \leq \|\partial_x \varphi(x)\|_\infty\}$$

is always equal to

$$K := \max\{|H(x, u, p)| : x \in M, |u| \leq l_{k-1}, \|p\| \leq \|\partial_x \varphi(x)\|_\infty\}.$$

Note that $u_{n,k}$ is the solution of

$$\partial_t u + H_n(x, u_{n,k-1}(x, t), \partial_x u) = 0. \quad (2.11)$$

Similar to the proof of (2.3), we have

$$\partial_t u_{n,k}(x, t) \geq -K_n - \lambda \|\partial_t u_{n,k-1}\|_\infty T.$$

Combining (2.11) and the definition of H_n , we have

$$\begin{aligned} H(x, 0, \partial_x u_{n,k}(x, t)) &\leq H_n(x, 0, \partial_x u_{n,k}(x, t)) \\ &\leq K + \lambda \|\partial_t u_{n,k-1}\|_\infty T + \lambda l_{k-1}, \quad \forall n \geq \|\partial_x \varphi(x)\|_\infty. \end{aligned}$$

Therefore, $\{u_{n,k}(x, t)\}_{n \in \mathbb{N}}$ is equi-Lipschitz. Note that

$$u_{n,k}(x, 0) = \varphi(x).$$

It follows that $\{u_{n,k}(x, t)\}_{n \in \mathbb{N}}$ is uniformly bounded, so it has a converging subsequence. We have to show that all converging subsequences have the same limit function u_k . In fact, according to Lemma A.7, the value function

$$\bar{u}_{n,k}(x, t) := \inf_{\gamma(t)=x} \left\{ \varphi(\gamma(0)) + \int_0^t L_n(\gamma(\tau), u_{k-1}(\gamma(\tau), \tau), \dot{\gamma}(\tau)) d\tau \right\}$$

converges to $u_k(x, t)$ pointwisely. Taking a minimizer γ of $u_{n,k}(x, t)$, we have

$$\begin{aligned} \bar{u}_{n,k}(x, t) - u_{n,k}(x, t) &\leq \varphi(\gamma(0)) + \int_0^t L_n(\gamma(\tau), u_{k-1}(\gamma(\tau), \tau), \dot{\gamma}(\tau)) d\tau \\ &\quad - \varphi(\gamma(0)) + \int_0^t L_n(\gamma(\tau), u_{n,k-1}(\gamma(\tau), \tau), \dot{\gamma}(\tau)) d\tau \\ &\leq \lambda \|u_{k-1}(x, t) - u_{n,k-1}(x, t)\|_\infty T. \end{aligned}$$

Exchanging the role of $\bar{u}_{n,k}(x, t)$ and $u_{n,k}(x, t)$, we have $\|\bar{u}_{n,k}(x, t) - u_{n,k}(x, t)\|_\infty \rightarrow 0$ as $n \rightarrow \infty$. It follows that

$$\lim_{n \rightarrow +\infty} u_{n,k}(x, t) = u_k(x, t), \quad \text{uniformly on } M \times [0, T],$$

which implies that $u_k(x, t)$ is Lipschitz. Note that the Lipschitz constant may depend on k . Thus, the assertion A[k] holds.

Since H_n converges uniformly to H on compact subsets of $T^*M \times \mathbb{R}$, and $u_{n,k}(x, t)$ converges uniformly to $u_k(x, t)$ on $M \times [0, T]$, by the stability of the viscosity solutions, we conclude that $u_k(x, t)$ is the Lipschitz solution of (2.2). \square

By Lemma 2.2 (i), $u_k(x, t)$ converges uniformly to $u(x, t)$ on $M \times [0, T]$. Moreover, $\sup_{k \in \mathbb{N}} \|u_k(x, t)\|_\infty$ is finite. Since $\varphi \in \text{Lip}(M)$, then $\|\partial_x \varphi\|_\infty$ is also finite. By Lemma 2.2 (ii), $\{u_k(x, t)\}_{k \in \mathbb{N}}$ is equi-Lipschitz. Therefore the limit function $u(x, t) := T_t^- \varphi(x)$ is the Lipschitz solution of (CP_H) . Theorem 1 has been proved when φ is Lipschitz.

2.2 On Item (**): the continuous initial conditions

At the beginning, we need to show $T_t^- \varphi$ is well-defined for each $\varphi \in C(M)$. By Lemma 2.2, it suffices to prove that given $T > 0$ and $\varphi \in C(M)$, u_k defined in (2.1) is continuous on $M \times [0, T]$.

In fact, for any $\varphi \in C(M)$, there exists a sequence of Lipschitz functions $\{\varphi_m\}_{m \in \mathbb{N}}$ converging uniformly to φ . We have already proved in Lemma 2.4 that for the initial function φ_m , the solution of (2.2), denoted by $u_k^m(x, t)$, is Lipschitz. We then proceed by induction in the following. By definition, u_0^m converges uniformly to u_0 . Assume u_{k-1}^m converges uniformly to u_{k-1} , then u_{k-1} is continuous. By Lemma 2.1(i), $u_k(x, t)$ admits a minimizer γ . By definition, we have

$$u_k^m(x, t) - u_k(x, t) \leq \varphi_m(\gamma(0)) - \varphi(\gamma(0)) + \lambda \|u_{k-1}^m(x, t) - u_{k-1}(x, t)\|_\infty T.$$

Exchanging the roles of $u_k^m(x, t)$ and $u_k(x, t)$, we obtain $\|u_k^m - u_k\|_\infty \rightarrow 0$ as $m \rightarrow \infty$. Therefore, u_k defined in (2.1) is continuous on $M \times [0, T]$.

By Lemma 2.2 (i), $u_k(x, t)$ converges uniformly to $u(x, t) := T_t^- \varphi(x)$. It follows that $u(x, t)$ is continuous. It remains to verify that $u(x, t)$ is the solution of (CP_H) .

We have proved in Item (*) that for $\varphi \in \text{Lip}(M)$, $T_t^- \varphi(x)$ is the Lipschitz solution of (CP_H) . We assert for any φ and $\psi \in C(M)$,

$$\|T_t^- \varphi - T_t^- \psi\|_\infty \leq e^{\lambda t} \|\varphi - \psi\|_\infty. \quad (2.12)$$

If the assertion is true, then for $t \in [0, T]$, $T_t^- \varphi_m$ converges uniformly to $T_t^- \varphi$ as $m \rightarrow \infty$. According to the stability of viscosity solutions, we conclude that $T_t^- \varphi$ is the solution of (CP_H) under the initial condition $u(x, 0) = \varphi(x)$. The uniqueness of the solution of (CP_H) is guaranteed by the comparison theorem (see [18, Theorem 2.1]). The assertion (2.12) above will be verified in Proposition 3.1 below.

3 An existence result for the solutions of (E_H)

In order to prove Theorem 2, we collect two basic properties of the backward and forward Lax-Oleinik semigroups in the following.

Proposition 3.1.

- (1) For any φ and $\psi \in C(M)$, if $\varphi(x) < \psi(x)$ for all $x \in M$, then $T_t^- \varphi(x) < T_t^- \psi(x)$ and $T_t^+ \varphi(x) < T_t^+ \psi(x)$ for all $(x, t) \in M \times (0, +\infty)$.

(2) For any φ and $\psi \in C(M)$, then $\|T_t^- \varphi - T_t^- \psi\|_\infty \leq e^{\lambda t} \|\varphi - \psi\|_\infty$ and $\|T_t^+ \varphi - T_t^+ \psi\|_\infty \leq e^{\lambda t} \|\varphi - \psi\|_\infty$ for all $t > 0$.

Proof. We first prove Item (1). Assume that there exists $(x, t) \in M \times [0, +\infty)$ such that $T_t^- \varphi(x) \geq T_t^- \psi(x)$. Let $\gamma : [0, t] \rightarrow M$ be a minimizer of $T_t^- \psi(x)$ with $\gamma(t) = x$. Define

$$F(s) = T_s^- \psi(\gamma(s)) - T_s^- \varphi(\gamma(s)), \quad s \in [0, t].$$

Then F is a continuous function defined on $[0, t]$, and $F(0) > 0$. By assumption we have $F(t) \leq 0$. Then there is $s_0 \in [0, t)$ such that $F(s_0) = 0$ and $F(s) > 0$ for all $s \in [0, s_0)$. Since γ is a minimizer of $T_t^- \varphi_2(x)$, we have

$$T_{s_0}^- \psi(\gamma(s_0)) = T_{s_0}^- \psi(\gamma(s_0)) + \int_s^{s_0} L(\gamma(\tau), T_\tau^- \psi(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau,$$

and

$$T_{s_0}^- \varphi(\gamma(s_0)) \leq T_{s_0}^- \varphi(\gamma(s_0)) + \int_s^{s_0} L(\gamma(\tau), T_\tau^- \varphi(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau,$$

which implies $F(s_0) \geq F(s) - \lambda \int_s^{s_0} F(\tau) d\tau$. Here $F(s_0) = 0$, thus

$$F(s) \leq \lambda \int_s^{s_0} F(\tau) d\tau.$$

By the Gronwall inequality, we conclude $F(s) \equiv 0$ for all $s \in [0, s_0)$, which contradicts $F(0) > 0$.

Next, we prove Item (2). For a given $x \in M$ and $t > 0$, if $T_t^- \varphi(x) = T_t^- \psi(x)$, then the proof is completed. Without loss of generality, we consider $T_t^- \varphi(x) > T_t^- \psi(x)$. Let $\gamma : [0, t] \rightarrow M$ be a minimizer of $T_t^- \psi(x)$, define

$$F(s) := T_s^- \varphi(\gamma(s)) - T_s^- \psi(\gamma(s)), \quad \forall s \in [0, t].$$

By assumption we have $F(t) > 0$. If there is $\sigma \in [0, t)$ such that $F(\sigma) = 0$ and $F(s) > 0$ for all $s \in (\sigma, t]$, by definition we have

$$T_s^- \varphi(\gamma(s)) \leq T_t^- \varphi(\gamma(\sigma)) + \int_\sigma^s L(\gamma(\tau), T_\tau^- \varphi(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau,$$

and

$$T_s^- \psi(\gamma(s)) = T_t^- \psi(\gamma(\sigma)) + \int_\sigma^s L(\gamma(\tau), T_\tau^- \psi(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau,$$

which implies

$$F(s) \leq F(\sigma) + \lambda \int_\sigma^s F(\tau) d\tau,$$

where $F(\sigma) = 0$. By the Gronwall inequality we conclude $F(s) \equiv 0$ for all $s \in [\sigma, t]$, which contradicts $F(t) > 0$.

Therefore, for all $\sigma \in [0, t]$, we have $F(\sigma) > 0$. Here $0 < F(0) \leq \|\varphi - \psi\|_\infty$. By definition we have

$$T_s^- \varphi(\gamma(\sigma)) \leq T_t^- \varphi(\gamma(0)) + \int_0^\sigma L(\gamma(\tau), T_\tau^- \varphi(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau,$$

and

$$T_s^- \psi(\gamma(\sigma)) = T_t^- \psi(\gamma(0)) + \int_0^\sigma L(\gamma(\tau), T_\tau^- \psi(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau,$$

which implies

$$F(\sigma) \leq F(0) + \lambda \int_0^\sigma F(\tau) d\tau.$$

By the Gronwall inequality we get $F(\sigma) \leq \|\varphi - \psi\|_\infty e^{\lambda\sigma}$, which implies $T_t^- \varphi(x) - T_t^- \psi(x) \leq \|\varphi - \psi\|_\infty e^{\lambda t}$ by taking $\sigma = t$. Exchanging the role of φ and ψ , we finally obtain that $\|T_t^- \varphi(x) - T_t^- \psi(x)\|_\infty \leq \|\varphi - \psi\|_\infty e^{\lambda t}$.

By definition, one can show the corresponding properties of T^+ . \square

Generally speaking, the local boundedness of $L(x, u, \dot{x})$ does not hold if $H(x, u, p)$ satisfies the assumption (CER) rather than superlinearity. Fortunately, similar to [19, Lemma 2.3], one can prove the local boundedness of $L(x, u, \dot{x})$ restricting on certain regions.

Lemma 3.2. *Let $H(x, 0, p)$ satisfy (C)(CON)(CER), there exist constants $\delta > 0$ and $\bar{C} > 0$ such that the Lagrangian $L(x, 0, \dot{x})$ associated to $H(x, 0, p)$ satisfies*

$$L(x, 0, \xi) \leq \bar{C}, \quad \forall (x, \xi) \in M \times \bar{B}(0, \delta).$$

In the following part of this paper, we define

$$\mu := \frac{\text{diam}(M)}{\delta}. \quad (3.1)$$

Lemma 3.3. *Let $\varphi \in C(M)$.*

- (1) *Given any $x_0 \in M$, if $T_t^- \varphi(x_0)$ does not have an upper bound as $t \rightarrow +\infty$, then for any $c \in \mathbb{R}$, there exists $t_c > 0$ such that $T_{t_c}^- \varphi(x) > \varphi(x) + c$ for all $x \in M$.*
- (2) *Given any $x_0 \in M$, if $T_t^- \varphi(x_0)$ does not have a lower bound as $t \rightarrow +\infty$, then for any $c \in \mathbb{R}$, there exists $t_c > 0$ such that $T_{t_c}^- \varphi(x) < \varphi(x) + c$ for all $x \in M$.*

Proof. We only prove Item (1). Item (2) is similar to be verified. Assume that there exists $c_0 \in \mathbb{R}$ such that for any $t > 0$, we have a point $x_t \in M$ satisfying $T_t^- \varphi(x_t) \leq \varphi(x_t) + c_0$. Let $\alpha : [0, \mu] \rightarrow M$ be a geodesic connecting x_t and x with constant speed, where the constant μ was defined in (3.1), then $\|\dot{\alpha}\| \leq \delta$. If $T_{t+\mu}^- \varphi(x) > \varphi(x) + c_0$, since $T_t^- \varphi(x_t) \leq \varphi(x_t) + c_0$, there exists $\sigma \in [0, \mu]$ such that $T_{t+\sigma}^- \varphi(\alpha(\sigma)) = \varphi(x_t) + c_0$ and $T_{t+s}^- \varphi(\alpha(s)) > \varphi(x_t) + c_0$ for all $s \in (\sigma, \mu]$. By definition we have

$$\begin{aligned} T_{t+s}^- \varphi(\alpha(s)) &\leq T_{t+\sigma}^- \varphi(\alpha(\sigma)) + \int_\sigma^s L(\alpha(\tau), T_{t+\tau}^- \varphi(\alpha(\tau)), \dot{\alpha}(\tau)) d\tau \\ &= \varphi(x_t) + c_0 + \int_\sigma^s L(\alpha(\tau), T_{t+\tau}^- \varphi(\alpha(\tau)), \dot{\alpha}(\tau)) d\tau, \end{aligned}$$

which implies

$$\begin{aligned} T_{t+s}^- \varphi(\alpha(s)) - (\varphi(x_t) + c_0) &\leq \int_\sigma^s L(\alpha(\tau), T_{t+\tau}^- \varphi(\alpha(\tau)), \dot{\alpha}(\tau)) d\tau \\ &\leq \int_\sigma^s L(\alpha(\tau), \varphi(x_t) + c_0, \dot{\alpha}(\tau)) d\tau + \lambda \int_\sigma^s (T_{t+\tau}^- \varphi(\alpha(\tau)) - (\varphi(x_t) + c_0)) d\tau \\ &\leq L_0 \mu + \lambda \int_\sigma^s (T_{t+\tau}^- \varphi(\alpha(\tau)) - (\varphi(x_t) + c_0)) d\tau, \end{aligned}$$

where

$$L_0 := \bar{C} + \lambda \|\varphi + c_0\|_\infty,$$

and \bar{C} is given in Lemma 3.2. By the Gronwall inequality, we have

$$T_{t+s}^- \varphi(\alpha(s)) - (\varphi(x_t) + c_0) \leq L_0 \mu e^{\lambda(s-\sigma)} \leq L_0 \mu e^{\lambda\mu}, \quad \forall s \in (\sigma, \mu].$$

Take $s = \mu$. We have $T_{t+\mu}^- \varphi(x) \leq \varphi(x_t) + c_0 + L_0 \mu e^{\lambda\mu}$. It means that $T_{t+\mu}^- \varphi(x)$ has an upper bound independent of t , which contradicts the assumption. \square

Lemma 3.4. *If there exist two continuous functions φ_1 and φ_2 on M such that*

$$T_t^- \varphi_1 \geq C_1, \quad T_t^- \varphi_2 \leq C_2,$$

then there is a constant function $\bar{\varphi}$ such that $|T_t^- \bar{\varphi}| \leq C_3$ for all $(x, t) \in M \times [0, +\infty)$, where C_i , $i = 1, 2, 3$, are constants independent of x and t .

Proof. Define $A_1 := \|\varphi_1\|_\infty$ and $A_2 := -\|\varphi_2\|_\infty$, then $A_2 \leq A_1$ and

$$T_t^- A_1(x) \geq T_t^- \varphi_1(x), \quad T_t^- A_2(x) \leq T_t^- \varphi_2(x) \quad \text{for all } x \in M.$$

If $T_t^- A_1(x)$ has an upper bound independent of t , then $\bar{\varphi} \equiv A_1$ is enough. If $T_t^- A_1(x)$ does not have an upper bound independent of t , we define

$$A^* := \inf\{A : \exists t_A > 0 \text{ such that } T_{t_A}^- A(x) \geq A, \forall x \in M\}.$$

By using Lemma 3.3 (1) with $c = 0$, we have $A^* \leq A_1 < +\infty$. The remaining discussion is divided into two cases.

Case (1): $A^* > -\infty$. In this case, we aim to prove that $\bar{\varphi} \equiv A^*$ is enough.

We first show that $T_t^- A^*(x)$ has an upper bound independent of t . Assume $T_t^- A^*(x)$ does not have an upper bound. By Lemma 3.3 (1), for $c = 1$, there is $t_1 > 0$ such that $T_{t_1}^- A^*(x) > A^* + 1$ for all $x \in M$. By Proposition 3.1 (2), for any $\varepsilon > 0$, we have

$$T_{t_1}^- (A^* - \varepsilon)(x) \geq T_{t_1}^- A^*(x) - e^{\lambda t_1} \varepsilon > A^* + 1 - e^{\lambda t_1} \varepsilon.$$

For every $0 < \varepsilon < (e^{\lambda t_1} - 1)^{-1}$, we have $T_{t_1}^- (A^* - \varepsilon)(x) > A^* - \varepsilon$. It means that we have found a smaller constant $A^* - \varepsilon$ such that if $t_{A^* - \varepsilon} := t_1$, then

$$T_{t_{A^* - \varepsilon}}^- (A^* - \varepsilon)(x) > A^* - \varepsilon,$$

which contradicts the definition of A^* .

We then prove that $T_t^- A^*$ has a lower bound independent of t . Assume $T_t^- A^*(x)$ does not have a lower bound. By using Lemma 3.3 (2) with $c = -1$, there is $t_1 > 0$ such that $T_{t_1}^- A^*(x) < A^* - 1$ for all $x \in M$. Since $T_t^- A^*(x)$ has an upper bound independent of t , then $A^* < A_1$. By Proposition 3.1 (2) and $A^* < A_1$, there is a constant $\delta_0 > 0$ such that $A^* + \delta < A_1$ and

$$T_{t_1}^- (A^* + \delta)(x) < A^* - \frac{1}{2} + \delta < A^* + \delta, \quad (3.2)$$

for all $\delta \in [0, \delta_0)$. By the definition of A^* , there is $\bar{A} \in [A^*, A^* + \delta_0)$ and $t_2 := t_{\bar{A}} > 0$ such that

$$T_{t_2}^- \bar{A}(x) \geq \bar{A}. \quad (3.3)$$

By (3.2), we have

$$T_{t_1}^- \bar{A}(x) < \bar{A} - \frac{1}{2} < \bar{A}. \quad (3.4)$$

Define $B^* := \bar{A} - \frac{1}{2}$. According to the continuity of $T_t^- \varphi(x)$ at $t = 0$, there exists $\varepsilon_0 > 0$ such that for $0 \leq \sigma < \varepsilon_0$, we have

$$T_\sigma^- B^*(x) \leq \bar{A} - \frac{1}{4}. \quad (3.5)$$

For t_1 and $t_2 > 0$, there exist n_1 and $n_2 \in \mathbb{N}$, and $\varepsilon \in [0, \varepsilon_0)$ such that $n_1 t_1 + \varepsilon = n_2 t_2$. By Proposition 3.1 (1) and (3.2), we have

$$T_{n_1 t_1}^- \bar{A}(x) \leq T_{t_1}^- \bar{A}(x) < B^*. \quad (3.6)$$

Take $\sigma = \varepsilon$ in (3.5). By Proposition 3.1 (1) and (3.6), we get

$$T_\varepsilon^- \circ T_{n_1 t_1}^- \bar{A}(x) \leq T_\varepsilon^- B^*(x) \leq \bar{A} - \frac{1}{4}. \quad (3.7)$$

By (3.3), one has $T_{n_2 t_2}^- \bar{A}(x) \geq \bar{A}$. Thus

$$\bar{A} - \frac{1}{4} \geq T_\varepsilon^- \circ T_{n_1 t_1}^- \bar{A}(x) = T_{n_2 t_2}^- \bar{A}(x) \geq \bar{A}, \quad (3.8)$$

which is a contradiction.

Case (2): $A^* = -\infty$. In this case, we aim to prove that for any $A < A_2$, the function $T_t^- A(x)$ is uniformly bounded. Namely, $\bar{\varphi} \equiv A$ is enough. Since $T_t^- A(x) \leq T_t^- A_2(x)$, then $T_t^- A(x)$ has an upper bound. The proof of the existence of the lower bound of $T_t^- A(x)$ is similar to Case (1). In fact, we only need to replace A^* , A_1 by A and A_2 respectively. \square

Remark 3.5. Let $\varphi \in C(M)$. According to [18, Theorem 6.1], if $T_t^- \varphi(x)$ has a bound independent of t , then the lower half limit

$$\check{\varphi}(x) := \lim_{r \rightarrow 0^+} \inf \{ T_t^- \varphi(y) : d(x, y) < r, t > 1/r \}$$

is a Lipschitz solution of (E_H) . According to Proposition D.4, the function $\check{\varphi}$ is a backward weak KAM solution of (E_H) . Similarly, if $T_t^+ \varphi(x)$ has a bound independent of t , define

$$\begin{aligned} \hat{\varphi}(x) &:= \lim_{r \rightarrow 0^+} \sup \{ T_t^+ \varphi(y) : d(x, y) < r, t > 1/r \} \\ &= \lim_{r \rightarrow 0^+} \sup \{ -\bar{T}_t^- (-\varphi)(y) : d(x, y) < r, t > 1/r \} \\ &= - \lim_{r \rightarrow 0^+} \inf \{ \bar{T}_t^- (-\varphi)(y) : d(x, y) < r, t > 1/r \}. \end{aligned}$$

Then $-\hat{\varphi}$ is a Lipschitz solution of $H(x, -u, -\partial_x u) = 0$. Equivalently, $\hat{\varphi}$ is a forward weak KAM solution of (E_H) .

Proof of Theorem 2. By assumption, there is $\varphi \in C(M)$ and $t_1 > 0$ such that $T_{t_1}^- \varphi \geq \varphi$. For any $t > 0$, one can find $n \in \mathbb{N}$ and $r \in [0, t_1)$ such that $t = nt_1 + r$. By Proposition 3.1 (1), we have $T_t^- \varphi \geq T_r^- \varphi$. Namely, $T_t^- \varphi$ has a lower bound independent of t . On the other hand, there is $\psi \in C(M)$ and $t_2 > 0$ such that $T_{t_2}^- \psi \leq \psi$. It is similar to obtain that $T_t^- \psi$ has an upper bound independent of t . By Lemma 3.4, there exists a constant function $\bar{\varphi}$ such that $T_t^- \bar{\varphi}$ is uniformly bounded. By Remark 3.5, (E_H) admits Lipschitz viscosity solutions. \square

4 The Aubry set

Let $u_- \in \mathcal{S}_-$. At the beginning, we prove that the limit function $x \mapsto \lim_{t \rightarrow +\infty} T_t^+ u_-(x)$ is well defined. Corollaries 4.3 and 4.7 guarantee the boundedness of $T_t^+ u_-$. Moreover, Item (1) of Theorem 3 is verified by Proposition 4.8, and Item (2) is shown by Proposition 4.9.

Proposition 4.1. *Let $\varphi \in C(M)$ and $u_- \in \mathcal{S}_-$. If φ satisfies the following condition:*

(\odot) $\varphi \leq u_-$ and there exists a point x_0 such that $\varphi(x_0) = u_-(x_0)$.

then $T_t^+ \varphi(x)$ has a bound independent of t and φ .

We divide the proof into three parts, that is, Lemmas 4.2, 4.5 and 4.6.

Lemma 4.2. *Suppose φ satisfies the condition (\odot), then $T_t^+ \varphi(x) \leq u_-(x)$ for all $t > 0$.*

Proof. Assume there exists $(x, t) \in M \times (0, +\infty)$ such that $T_t^+ \varphi(x) > u_-(x)$. Let $\gamma : [0, t] \rightarrow M$ be a minimizer of $T_t^+ \varphi(x)$ with $\gamma(0) = x$. Define

$$F(s) = T_{t-s}^+ \varphi(\gamma(s)) - u_-(\gamma(s)), \quad s \in [0, t].$$

Then $F(s)$ is continuous and $F(t) = \varphi(\gamma(t)) - u_-(\gamma(t)) \leq 0$. By assumption we have $F(0) > 0$. Then there is $\tau_0 \in (0, t]$ such that $F(\tau_0) = 0$ and $F(\tau) > 0$ for all $s \in [0, \tau_0)$. For each $\tau \in [0, \tau_0]$, we have

$$T_{t-\tau}^+ \varphi(\gamma(\tau)) = T_{t-\tau_0}^+ \varphi(\gamma(\tau_0)) - \int_{\tau}^{\tau_0} L(\gamma(s), T_{t-s}^+ \varphi(\gamma(s)), \dot{\gamma}(s)) ds.$$

Since $u_- = T_t^- u_-$ for all $t > 0$, we have

$$u_-(\gamma(\tau_0)) \leq u_-(\gamma(\tau)) + \int_{\tau}^{\tau_0} L(\gamma(s), u_-(\gamma(s)), \dot{\gamma}(s)) ds.$$

Thus $F(\tau) \leq F(\tau_0) + \lambda \int_{\tau}^{\tau_0} F(s) ds$, where $F(\tau_0) = 0$. Define $F(s) = G(\tau_0 - s)$. We get

$$G(\tau_0 - \tau) \leq \lambda \int_0^{\tau_0 - \tau} G(\sigma) d\sigma.$$

By the Gronwall inequality, we conclude $F(\tau) = G(\tau_0 - \tau) \equiv 0$ for all $\tau \in [0, \tau_0]$, which contradicts $F(0) > 0$. \square

Corollary 4.3. *Let $u_- \in \mathcal{S}_-$. Then $T_t^+ u_- \leq u_-$ for each $t > 0$.*

Combining Corollary 4.3 with Proposition 3.1 (1), one can obtain that $T_t^+ u_- = T_s^+ \circ T_{t-s}^+ u_- \leq T_s^+ u_-$ for all $t > s$, then we have

Corollary 4.4. *$T_t^+ u_-$ is decreasing in t .*

Lemma 4.5. *Suppose φ satisfies the condition (\odot). Let $\gamma_- : (-\infty, 0] \rightarrow M$ be a $(u_-, L, 0)$ -calibrated curve with $\gamma_-(0) = x_0$, then $T_t^+ \varphi(\gamma_-(-t)) = u_-(\gamma_-(-t))$ for each $t > 0$.*

Proof. For each $t > 0$, we define $\gamma_t(s) := \gamma_-(s - t)$ for $s \in [0, t]$. By Lemma 4.2, for each $s \in [0, t]$, we have $u_-(\gamma_t(s)) \geq T_{t-s}^+ \varphi(\gamma_t(s))$. Define

$$F(s) = u_-(\gamma_t(s)) - T_{t-s}^+ \varphi(\gamma_t(s)),$$

then $F(s) \geq 0$ and $F(t) = 0$. If $F(0) > 0$, then there is $s_0 \in (0, t]$ such that $F(s_0) = 0$ and $F(s) > 0$ for all $s \in [0, s_0)$. By definition, for $s_1 \in [0, s_0)$, we have

$$u_-(\gamma_t(s_0)) - u_-(\gamma_t(s_1)) = \int_{s_1}^{s_0} L(\gamma_t(s), u_-(\gamma_t(s)), \dot{\gamma}_t(s)) ds,$$

and

$$T_{t-s_1}^+ \varphi(\gamma_t(s_1)) \geq T_{t-s_0}^+ \varphi(\gamma_t(s_0)) - \int_{s_1}^{s_0} L(\gamma_t(s), T_{t-s}^+ \varphi(\gamma_t(s)), \dot{\gamma}_t(s)) ds,$$

which implies

$$F(s_1) \leq F(s_0) + \lambda \int_{s_1}^{s_0} F(s) ds.$$

By the Gronwall inequality, we conclude $F(s) \equiv 0$ for all $s \in [0, s_0]$, which contracts $F(0) > 0$. Therefore $F(0) = 0$. Namely, $T_t^+ \varphi(\gamma_t(0)) = u_-(\gamma_t(0))$. Recall $\gamma_t(s) := \gamma_-(s - t)$. We have $T_t^+ \varphi(\gamma_-(-t)) = u_-(\gamma_-(-t))$. \square

Lemma 4.6. *Suppose φ satisfies the condition (\odot) , then $T_t^+ \varphi(x)$ has a lower bound independent of t and φ .*

Proof. Let $\gamma_- : (-\infty, 0] \rightarrow M$ be a $(u_-, L, 0)$ -calibrated curve with $\gamma_-(0) = x_0$. Let $t > \mu$ and $\alpha : [0, \mu] \rightarrow M$ be a geodesic connecting x and $\gamma_-(-t + \mu)$ with constant speed, then $\|\dot{\alpha}\| \leq \delta$. If $T_t^+ \varphi(x) \geq u_-(\gamma_-(-t + \mu))$, then the proof is completed. It remains to consider $T_t^+ \varphi(x) < u_-(\gamma_-(-t + \mu))$. Since

$$T_{t-\mu}^+ \varphi(\gamma_-(-t + \mu)) = u_-(\gamma_-(-t + \mu)),$$

then there is $\sigma \in (0, \mu]$ such that

$$T_{t-\sigma}^+ \varphi(\alpha(\sigma)) = u_-(\gamma_-(-t + \mu)), \quad T_{t-s}^+ \varphi(\alpha(s)) < u_-(\gamma_-(-t + \mu)) \quad \text{for all } s \in [0, \sigma).$$

By definition we have

$$\begin{aligned} T_{t-s}^+ \varphi(\alpha(s)) &\geq T_{t-\sigma}^+ \varphi(\alpha(\sigma)) - \int_s^\sigma L(\alpha(\tau), T_{t-\tau}^+ \varphi(\alpha(\tau)), \dot{\alpha}(\tau)) d\tau \\ &= u_-(\gamma_-(-t + \mu)) - \int_s^\sigma L(\alpha(\tau), T_{t-\tau}^+ \varphi(\alpha(\tau)), \dot{\alpha}(\tau)) d\tau, \end{aligned}$$

which implies

$$\begin{aligned} u_-(\gamma_-(-t + \mu)) - T_{t-s}^+ \varphi(\alpha(s)) &\leq \int_s^\sigma L(\alpha(\tau), T_{t-\tau}^+ \varphi(\alpha(\tau)), \dot{\alpha}(\tau)) d\tau \\ &\leq \int_s^\sigma L(\alpha(\tau), u_-(\gamma_-(-t + \mu)), \dot{\alpha}(\tau)) d\tau + \lambda \int_s^\sigma (u_-(\gamma_-(-t + \mu)) - T_{t-\tau}^+ \varphi(\alpha(\tau))) d\tau \\ &\leq L_0 \mu + \lambda \int_s^\sigma (u_-(\gamma_-(-t + \mu)) - T_{t-\tau}^+ \varphi(\alpha(\tau))) d\tau, \end{aligned}$$

where

$$L_0 := \bar{C} + \lambda \|u_-\|_\infty,$$

and \bar{C} is given by Lemma 3.2. Let $G(\sigma - s) = u_-(\gamma_-(-t + \mu)) - T_{t-s}^+ \varphi(\alpha(s))$, then

$$G(\sigma - s) \leq L_0 \mu + \lambda \int_0^{\sigma-s} G(\tau) d\tau.$$

By the Gronwall inequality, we have

$$u_-(\gamma_-(-t + \mu)) - T_{t-s}^+ \varphi(\alpha(s)) = G(\sigma - s) \leq L_0 \mu e^{\lambda(\sigma-s)} \leq L_0 \mu e^{\lambda \mu}, \quad \forall s \in [0, \sigma].$$

Thus $T_t^+ \varphi(x) \geq u_-(\gamma_-(-t + \mu)) - L_0 \mu e^{\lambda \mu}$. We finally get a lower bound of $T_t^+ \varphi(x)$ independent of t and φ . \square

Corollary 4.7. $T_t^+ u_-$ has a lower bound independent of t .

Proposition 4.8. $T_t^+ u_-$ converges to a forward weak KAM solution u_+ of (E_H) uniformly as $t \rightarrow +\infty$.

Proof. We first recall that $T_t^+ \varphi := -\bar{T}_t^-(-\varphi)$, where \bar{T}_t^- denotes the backward Lax-Oleinik semigroup associated to $L(x, -u, -\dot{x})$. Since $T_t^+ u_-$ is decreasing in t , the function $u(x, t) := \bar{T}_t^-(-u_-)$ is increasing in t . Thus, $\partial_t u(x, t) \geq 0$ holds in the viscosity sense. Since $u(x, t)$ is the viscosity solution of $\partial_t u + H(x, -u, -\partial_x u) = 0$, we have $H(x, -u, -\partial_x u) \leq 0$. Since $T_t^+ u_-$ has a bound independent of t , $u(x, t)$ has a bound independent of t . We conclude that $\|\partial_x T_t^+ u_-\|_\infty = \|\partial_x u(x, t)\|_\infty$ has a bound independent of t by (CER). Corollaries 4.4 and 4.7 imply that the pointwise limit $u_+(x) = \lim_{t \rightarrow +\infty} T_t^+ u_-(x)$ exists. Since $\|\partial_x T_t^+ u_-\|_\infty$ has a bound independent of t , the limit function u_+ is continuous. By the Dini theorem, the family $T_t^+ u_-$ converges uniformly to u_+ . It remains to prove that u_+ is a fixed point of T_t^+ . For each $t > 0$, by Proposition 3.1 (2), we have

$$\|T_{t+s}^+ u_- - T_t^+ u_+\|_\infty \leq e^{\lambda t} \|T_s^+ u_- - u_+\|_\infty.$$

Letting $s \rightarrow +\infty$, we get $T_t^+ u_+ = u_+$. \square

Proposition 4.9. The set \mathcal{I}_{u_-} is nonempty. More precisely, let $\gamma_- : (-\infty, 0] \rightarrow M$ be a $(u_-, L, 0)$ -calibrated curve. Define

$$\alpha(\gamma_-) := \{x \in M : \text{there exists a sequence } t_n \rightarrow -\infty \text{ such that } d(\gamma_-(t_n), x) \rightarrow 0\}.$$

Then $\alpha(\gamma_-)$ is nonempty, and it is contained in \mathcal{I}_{u_-} .

Proof. Let $\gamma_- : (-\infty, 0] \rightarrow M$ be a $(u_-, L, 0)$ -calibrated curve. By Lemma 4.5, for each $t > 0$ we have

$$T_t^+ u_-(\gamma_-(-t)) = u_-(\gamma_-(-t)).$$

Since M is compact, the set $\alpha(\gamma_-)$ is nonempty. Let $x^* \in \alpha(\gamma_-)$ and $t_n \rightarrow +\infty$ such that $d(\gamma_-(-t_n), x^*) \rightarrow 0$. The following inequality holds

$$\begin{aligned} |T_{t_n}^+ u_-(\gamma_-(-t_n)) - u_+(x^*)| &\leq |T_{t_n}^+ u_-(\gamma_-(-t_n)) - u_+(\gamma_-(-t_n))| \\ &\quad + |u_+(\gamma_-(-t_n)) - u_+(x^*)|. \end{aligned}$$

The function u_+ is Lipschitz (see Proposition D.2). Thus, as $t_n \rightarrow +\infty$,

$$|u_+(\gamma_-(-t_n)) - u_+(x^*)| \rightarrow 0.$$

Since $T_t^+ u_-$ converges to u_+ uniformly, then

$$|T_{t_n}^+ u_-(\gamma_-(-t_n)) - u_+(\gamma_-(-t_n))| \rightarrow 0.$$

Therefore, the limit of $T_{t_n}^+ u_-(\gamma_-(-t_n))$ is $u_+(x^*)$. On the other hand, we have

$$T_{t_n}^+ u_-(\gamma_-(-t_n)) = u_-(\gamma_-(-t_n)),$$

which tends to $u_-(x^*)$ by the continuity of u_- . We conclude that $u_+(x^*) = u_-(x^*)$. It means $\alpha(\gamma_-) \subseteq \mathcal{I}_{u_-}$. \square

5 A comparison result for the solutions of (E_H)

According to [8, Theorem 3.2], the viscosity solution of

$$H(x, -u(x), -\partial_x u(x)) = 0$$

is unique. By Proposition D.4, the forward weak KAM solution u_+ of (E_H) is also unique. Define $u_- = \lim_{t \rightarrow +\infty} T_t^- u_+$, then the conjugate pair (u_-, u_+) is unique. According to Proposition 4.8, $T_t^+ v_-$ converges to the unique forward weak KAM solution u_+ uniformly as $t \rightarrow +\infty$ and $u_+ \leq v_-$ for all $v_- \in \mathcal{S}_-$.

Proof of Theorem 4. We first prove the result (1). By Proposition 4.9, the set \mathcal{I}_{v_-} is nonempty for each $v_- \in \mathcal{S}_-$. For $x \in \mathcal{I}_{v_2}$, we have

$$u_+(x) \leq v_1(x) \leq v_2(x) = u_+(x),$$

then $v_1(x) = v_2(x) = u_+(x)$, that is, $x \in \mathcal{I}_{v_1}$.

We then prove the result (2). For each $x \in M$, let $\gamma_2 : (-\infty, 0] \rightarrow M$ be a $(v_2, L, 0)$ -calibrated curve with $\gamma_2(0) = x$. By Proposition 4.9, there is a $t_0 > 0$ large enough, such that $\gamma_2(-t_0) \in \mathcal{O}$, where \mathcal{O} denotes a neighborhood of \mathcal{I}_{v_2} . Define

$$F(s) = v_1(\gamma_2(s)) - v_2(\gamma_2(s)), \quad s \in [-t_0, 0].$$

If $v_1(x) > v_2(x)$, then $F(0) = v_1(x) - v_2(x) > 0$ and $F(-t_0) = v_1(\gamma_2(-t_0)) - v_2(\gamma_2(-t_0)) \leq 0$. Then there is $\sigma \in [-t_0, 0)$ such that $F(\sigma) = 0$ and $F(s) > 0$ for all $s \in (\sigma, 0]$. By definition we have

$$v_1(\gamma_2(s)) - v_1(\gamma_2(\sigma)) \leq \int_{\sigma}^s L(\gamma_2(\tau), v_1(\gamma_2(\tau)), \dot{\gamma}_2(\tau)) d\tau,$$

and

$$v_2(\gamma_2(s)) - v_2(\gamma_2(\sigma)) = \int_{\sigma}^s L(\gamma_2(\tau), v_2(\gamma_2(\tau)), \dot{\gamma}_2(\tau)) d\tau,$$

which implies

$$F(s) \leq F(\sigma) + \lambda \int_{\sigma}^s F(\tau) d\tau.$$

By the Gronwall inequality we conclude $F(s) \equiv 0$ for all $s \in [\sigma, 0]$, which contradicts $F(0) > 0$. We conclude $v_1 \leq v_2$ on M .

The result (3) follows directly from (2). The proof is now complete. \square

6 On the example (E1)

Let u_+ be the unique forward weak KAM solution of (E1). We have already known that $u_+ \leq v_-$ for each viscosity solution v_- of (E1). It is sufficient to show $u_+(x) < u_2(x)$ for all $x \in (-1, 1] \setminus \{0\}$. By the symmetry of u_2 , we only need to consider $x \in (0, 1]$.

By [5, Theorem 5.3.6] and Proposition D.4, each u_+ is a semiconvex function with linear modulus. Note that $u_+(x) \leq u_2(x)$. Moreover, u_+ can not be equal to u_2 at $x = 1$. In fact, if $u_2 = u_+$ at $x = 1$, combining with the semiconcavity of u_2 , then u_2 is differentiable at this point. Let us recall

$$u_2(x) = \frac{\lambda - \sqrt{\lambda^2 - 4}}{2} V(x),$$

and V is not differentiable at $x = 1$. This is a contradiction.

We then assume that there exists $x_0 \in (0, 1)$ such that $u_+(x_0) = u_2(x_0)$. Since $u_2(x)$ is differentiable for each $x \in [0, 1)$, then u_2 satisfies

$$-\lambda u(x) + \frac{1}{2}|u'(x)|^2 + V(x) = 0$$

in the classical sense for $x \in [0, 1)$. Note that $|u_2'(x)| > 0$ for $x \in (0, 1)$, we have $\lambda u_2(x) > V(x)$ for all $x \in (0, 1)$. For $z > V(x)$, we set

$$f(x, z) := \lambda \sqrt{2(z - V(x))},$$

then the function $(x, z) \mapsto f(x, z)$ is of class C^1 on

$$\{(x, z) \in \mathbb{R}^2 \mid x \in (0, 1), z > V(x)\}.$$

Given $\varepsilon \in (0, x_0)$, denote

$$\Omega_\varepsilon := \left\{ (x, z) \in \mathbb{R}^2 \mid x \in [\varepsilon, 1), z \in \left[\frac{1}{2}\lambda u_2(x) + \frac{1}{2}V(x), \frac{3}{2}\lambda u_2(x) - \frac{1}{2}V(x) \right] \right\}.$$

It follows that

$$\left| \frac{\partial f}{\partial z} \right| = \frac{\lambda}{\sqrt{2(z - V(x))}} \leq \frac{\lambda}{\sqrt{\lambda u_2(\varepsilon) - V(\varepsilon)}} < +\infty.$$

By the classical theory of ordinary differential equations, for $x_0 \in (0, 1)$, $\lambda u_2(x)$ is the unique solution of

$$\frac{dz}{dx} = f(x, z), \quad z(x_0) = \lambda u_2(x_0), \quad \text{on } \Omega_\varepsilon. \quad (6.1)$$

We assert that u_+ is differentiable on $(0, 1)$. If the assertion is true, then u_+ satisfies (E1) in the classical sense. Since $u_+ \leq u_2$ and $u_+(x_0) = u_2(x_0)$, λu_+ is the unique solution of (6.1) on Ω_ε . That is, $u_+ = u_2$ on $(\varepsilon, 1)$. Moreover, $u_+ = u_2$ on \mathbb{S} by continuity and the arbitrariness of ε . This contradicts the semiconvexity of u_+ . Therefore, we have $u_+(x) < u_2(x)$ for all $x \in (0, 1]$.

It remains to show that u_+ is differentiable on $(0, 1)$. Assume there exists $y_0 \in (0, 1)$ such that u_+ is not differentiable at y_0 . By [27, Lemma 2.2], [5, Theorem 3.3.6], and Proposition D.4, we have

$$D^*u_+(x) = \{p \in D^-u_+(x) \mid H(x, u_+(x), p) = 0\}, \quad D^-u_+(x) = \text{co}D^*u_+(x),$$

where D^* stands for the set of all reachable gradients and “co” denotes the convex hull. It follows from (E1) that

$$D^*u_+(y_0) = \{\pm l\},$$

where l is a positive constant. By the semiconvexity of u_+ , there exists $y_1 \in (0, y_0)$ such that $u_+(y_1) > u_+(y_0)$. Moreover, there is $z_0 \in (0, y_0)$ achieving a local maximum of u_+ . By using the semiconvexity of u_+ again, it is differentiable at z_0 , then $u'_+(z_0) = 0$. By (E1), we have

$$-\lambda u_+(z_0) + V(z_0) = 0.$$

Since $u'_+(x)$ exists for almost all x , there is $z_1 \in (z_0, y_0)$ such that $u'_+(z_1)$ exists. By the Newton-Leibniz formula, one can require $|u'_+(z_1)| > 0$ and $u_+(z_0) \geq u_+(z_1) \geq 0$. By definition, we have $V(z_1) > V(z_0)$. Therefore

$$-\lambda u_+(z_1) + \frac{1}{2}|u'_+(z_1)|^2 + V(z_1) > -\lambda u_+(z_0) + V(z_0) = 0,$$

which contradicts that u_+ satisfies (E1) at z_1 in the classical sense. \square

Acknowledgement The authors would like to thank Professor Wei Cheng and Kaizhi Wang for many useful discussions.

A One dimensional variational problems

The following results are useful in the proof of the existence and regularity of the minimizers in (T-), which all come from [4] and [22]. The results in [4, 22] were stated for the case in the Euclidean space \mathbb{R}^n . It is not difficult to generalize them for the case in the Riemannian manifold M .

A.1 Γ -convergence

Lemma A.1. *Let J be a bounded interval. Assume that $F(t, x, \dot{x})$ is lower semicontinuous, convex in \dot{x} , and has a lower bound. Then the integral functional*

$$\mathcal{F}(\gamma) = \int_J F(s, \gamma(s), \dot{\gamma}(s)) ds$$

is sequentially weakly lower semicontinuous in $W^{1,1}(J, M)$.

Proposition A.2. *Let M be a compact connected smooth manifold. Denote by $I = (a, b) \subset \mathbb{R}$ a bounded interval, and let $F(t, x, \dot{x})$ be a Lagrangian defined on $I \times TM$. Assume F satisfies*

- (i) $F(t, x, \dot{x})$ is measurable in t for all (x, \dot{x}) , and continuous in (x, \dot{x}) for almost every t ;
- (ii) $F(t, x, \dot{x})$ is convex in \dot{x} ;
- (iii) $F(t, x, \dot{x})$ is superlinear in \dot{x} .

Then for any given boundary condition x_0 and $x_1 \in M$, there exists a minimizer of $\int_I F(t, x, \dot{x}) dt$ in $\{x(t) \in W^{1,1}([a, b], M) : x(a) = x_0, x(b) = x_1\}$.

Definition A.3. Let X be a topological space. Given a sequence $F_n : X \rightarrow [-\infty, +\infty]$, then we define

$$\begin{aligned} (\Gamma - \liminf_{n \rightarrow +\infty} F_n)(x) &= \sup_{U \in \mathcal{N}(x)} \liminf_{n \rightarrow +\infty} \inf_{y \in U} F_n(y), \\ (\Gamma - \limsup_{n \rightarrow +\infty} F_n)(x) &= \sup_{U \in \mathcal{N}(x)} \limsup_{n \rightarrow +\infty} \inf_{y \in U} F_n(y). \end{aligned}$$

Here the neighbourhoods $\mathcal{N}(x)$ can be replaced by the topological basis. When the superior limit equals to the inferior limit, we can define the Γ -limit.

Definition A.4. Let X be a topological space. For every function $F : X \rightarrow [-\infty, +\infty]$, the lower semicontinuous envelope sc^-F of F is defined for every $x \in X$ by

$$(sc^-F)(x) = \sup_{G \in \mathcal{G}(F)} G(x),$$

where $\mathcal{G}(F)$ is the set of all lower semicontinuous functions G on X such that $G(y) \leq F(y)$ for every $y \in X$.

Lemma A.5. If F_n is an increasing sequence, then

$$\Gamma - \lim_{n \rightarrow +\infty} F_n = \lim_{n \rightarrow +\infty} sc^-F_n = \sup_{n \in \mathbb{N}} sc^-F_n.$$

Remark A.6. If F_n is an increasing sequence of lower semicontinuous functions which converges pointwisely to a function F , then F is lower semicontinuous and F_n has a Γ -convergence to F by Lemma A.5.

Lemma A.7. If the sequence F_n has a Γ -convergence in X to F , and there is a compact set $K \subset X$ such that

$$\inf_{x \in X} F_n(x) = \inf_{x \in K} F_n(x),$$

then F takes its minimum in X , and

$$\min_{x \in X} F(x) = \lim_{n \rightarrow +\infty} \inf_{x \in X} F_n(x).$$

A.2 Regularity of minimizers in t -dependent cases

The following results focus on the regularity of minimizers. Consider the following one dimensional variational problem

$$I(\gamma) := \int_a^b F(t, \gamma(t), \dot{\gamma}(t)) dt + \Psi(\gamma(a), \gamma(b)), \quad (\text{P})$$

where γ is taken in the class of absolutely continuous curves. Ψ takes its value in $\mathbb{R} \cup \{+\infty\}$ and stands for the constraints on the two ends of the curves γ .

In the following, we focus on a certain minimizer of the above integral functional, which is denoted by $\gamma_* \in W^{1,1}([a, b], M)$. Due to the Lavrentiev phenomenon, the minimizer may not be Lipschitz. One can refer [2] for various counterexamples. Thanks to [3], the Lipschitz regularity of the minimizers still holds for $F := L(x, v(x, t), \dot{x})$, where $v(x, t)$ is a Lipschitz function (see Lemma 2.3 (1)). Let us recall the related results in [3] as follows.

(\diamond): F takes its value in \mathbb{R} , there exist a constant $\varepsilon > 0$ and a Lebesgue-Borel-measurable map $k : [a, b] \times (0, +\infty) \rightarrow \mathbb{R}$ such that $k(t, 1) \in L^1[a, b]$, and, for a.e. $t \in [a, b]$, all $\sigma > 0$

$$|F(t_2, \gamma_*(t), \sigma \dot{\gamma}_*(t)) - F(t_1, \gamma_*(t), \sigma \dot{\gamma}_*(t))| \leq k(t, \sigma) |t_2 - t_1|,$$

where $t_1, t_2 \in [t - \varepsilon, t + \varepsilon] \cap [a, b]$.

Lemma A.8. *Let γ_* be a minimizer of (P). If F satisfies (\diamond), then there exists an absolutely continuous function $p \in W^{1,1}([a, b], \mathbb{R})$ such that for a.e. $t \in [a, b]$, we have*

$$F\left(t, \gamma_*(t), \frac{\dot{\gamma}_*(t)}{v}\right) v - F(t, \gamma_*(t), \dot{\gamma}_*(t)) \geq p(t)(v - 1), \quad \forall v > 0, \quad (\text{W})$$

and $|p'(t)| \leq k(t, 1)$ for a.e. $t \in [a, b]$.

Lemma A.9. *Let γ_* be a minimizer of (P). Assume F is a Borel measurable function. If F satisfies (\diamond) and*

(1) *Superlinearity: there exists a function $\Theta : [0, +\infty) \rightarrow \mathbb{R}$ satisfying*

$$\lim_{r \rightarrow +\infty} \frac{\Theta(r)}{r} = +\infty, \quad \text{and} \quad F(t, \gamma_*(t), \xi) \geq \Theta(\|\xi\|) \quad \text{for all } \xi \in T_{\gamma_*(t)}M.$$

(2) *Local boundedness: there exists $\rho > 0$ and $M \geq 0$ such that for a.e. $t \in [a, b]$, $F(t, \gamma_*(t), \xi) \leq M$ for all $\xi \in T_{\gamma_*(t)}M$ with $\|\xi\| = \rho$.*

Then the minimizer γ_* is Lipschitz. Moreover, if $\|\dot{\gamma}_*(t)\| > \rho$, we take $v = \|\dot{\gamma}_*(t)\|/\rho > 1$ in (W), then

$$F\left(t, \gamma_*(t), \rho \frac{\dot{\gamma}_*(t)}{\|\dot{\gamma}_*(t)\|}\right) \geq \rho \frac{\Theta(\|\dot{\gamma}_*(t)\|)}{\|\dot{\gamma}_*(t)\|} - \|p\|_\infty.$$

Therefore $\|\dot{\gamma}_*(t)\| \leq \max\{\rho, R\}$ where $R := \inf\{s : \rho \frac{\Theta(s)}{s} > M + \|p\|_\infty\}$.

B Proof of Lemma 2.1

When $H(x, u, p)$ is superlinear in p , it is well-known that the functional \mathbb{L}^t admits minimizers in $X_t(x)$. It remains to prove the existence of minimizers of \mathbb{L}^t when $H(x, u, p)$ is coercive in p . Define

$$\mathbb{L}_n^t(\gamma) = \varphi(\gamma(0)) + \int_0^t L_n(\gamma(s), v(\gamma(s), s), \dot{\gamma}(s)) ds,$$

where L_n is defined as in Section 2.1.2. Then each \mathbb{L}_n^t admits minimizers in $X_t(x)$. To prove the existence of the minimizers of $\mathbb{L}^t(\gamma)$, we define

$$m(r) := \inf_{x \in M} \left(\inf_{\|\dot{x}\| \geq r} L_1(x, 0, \dot{x}) \right), \quad \forall r \geq 0.$$

It is clear that the function $m(r)$ is superlinear, and

$$\begin{aligned} m(\|\dot{x}\|) &\leq L_n(x, 0, \dot{x}) \leq L_n(x, u, \dot{x}) + \lambda|u| \\ &\leq L(x, u, \dot{x}) + \lambda|u|, \quad \forall n \in \mathbb{N}, \quad \forall (x, u, \dot{x}) \in TM \times \mathbb{R}. \end{aligned}$$

For any sequence γ_n in $X_t(x)$ with $\lim_n \mathbb{L}^t(\gamma_n) < +\infty$, we have $\sup_n \int_0^t m(\|\dot{\gamma}_n\|)ds < +\infty$, so γ_n admits a weakly sequentially converging subsequence. By Lemma A.1, the functionals \mathbb{L}^t and \mathbb{L}_n^t are sequentially weakly lower semicontinuous on $X_t(x)$. Since $X_t(x)$ is a metric space, the functionals \mathbb{L}^t and \mathbb{L}_n^t are also lower semicontinuous. Note that $\{\mathbb{L}_n^t\}_{n \in \mathbb{N}}$ is an increasing sequence, and converges pointwisely to \mathbb{L}^t on $X_t(x)$. Both \mathbb{L}^t and $\mathbb{L}_n^t(\gamma)$ are lower semicontinuous. We conclude that $\Gamma - \lim_{n \rightarrow +\infty} \mathbb{L}_n^t = \mathbb{L}^t$ on $X_t(x)$ by Lemma A.5.

If the minimizers γ_n of \mathbb{L}_n^t are contained in a compact subset of $X_t(x)$, then by Lemma A.7, one can obtain that \mathbb{L}^t admits a minimum point on $X_t(x)$. It remains to show that there exists a compact set in $X_t(x)$ such that all minimizers γ_n are contained in this set. Consider the set

$$K_t(x) := \left\{ \gamma \in X_t(x) : \int_0^t m(\|\dot{\gamma}\|)ds \leq \|\phi\|_\infty + \mathbb{K}t + 2\lambda Kt \right\},$$

where $\mathbb{K} := \sup_{x \in M} L(x, 0, 0)$ and $K := \|v(x, t)\|_\infty$. The set $K_t(x)$ is weakly sequentially compact in $W^{1,1}([0, t], M)$. According to [4, Theorem 2.13], $K_t(x)$ is compact in $X_t(x)$. For the constant curve $\gamma_x \equiv x$, we have

$$\int_0^t m(\|\dot{\gamma}_x\|)ds \leq \mathbb{L}_n^t(\gamma_x) + \lambda Kt \leq \mathbb{L}^t(\gamma_x) + \lambda Kt \leq \|\phi\|_\infty + \mathbb{K}t + 2\lambda Kt.$$

Therefore γ_x is contained in $K_t(x)$. Similarly, for minimizers γ_n , we have

$$\begin{aligned} \int_0^t m(\|\dot{\gamma}_n\|)ds &\leq \mathbb{L}_n^t(\gamma_n) + \lambda Kt \leq \mathbb{L}_n^t(\gamma_x) + \lambda Kt \\ &\leq \mathbb{L}^t(\gamma_x) + \lambda Kt \leq \|\phi\|_\infty + \mathbb{K}t + 2\lambda Kt. \end{aligned}$$

Thus, all γ_n are contained in $K_t(x)$. □

C Proof of Lemma 2.3

Proof. We first prove Item (1). According to (LIP) and the Lipschitz continuity of $v(x, t)$ on $M \times [0, T]$, for each $\tau \in [0, t]$, the map $s \mapsto L(\gamma(\tau), v(\gamma(\tau), s), \dot{\gamma}(\tau))$ satisfies the condition (\diamond) , where $k \equiv \lambda \|\partial_t v(x, t)\|_\infty$. By Lemma A.9, for every $(x, t) \in M \times [0, T]$, the minimizers of $u(x, t)$ are Lipschitz. However, the Lipschitz constant depends on the end point (x, t) . We aim to show that for (x', t') sufficiently close to (x, t) , the Lipschitz constant of the minimizers of $u(x', t')$ is independent of (x', t') .

For any $r > 0$, if $d(x, x') \leq r$ and $|t - t'| \leq r/2$, where $t \geq r > 0$, we denote by $\gamma(s; x, t)$ and $\gamma(s; x', t')$ the minimizers of $u(x, t)$ and $u(x', t')$ respectively. Then we have

$$\begin{aligned} u(x', t') &= \varphi(\gamma(0; x', t')) + \int_0^{t'} L(\gamma(s; x', t'), v(\gamma(s; x', t'), s), \dot{\gamma}(s; x', t'))ds \\ &\leq \varphi(\gamma(0; x, t)) + \int_0^{t-r} L(\gamma(s; x, t), v(\gamma(s; x, t), s), \dot{\gamma}(s; x, t))ds \\ &\quad + \int_{t-r}^{t'} L(\alpha(s), v(\alpha(s), s), \dot{\alpha}(s))ds, \end{aligned}$$

where $\alpha : [t - r, t'] \rightarrow M$ is a geodesic connecting $\gamma(t - r; x, t)$ and x' with constant speed. Noticing that

$$\|\dot{\alpha}\| \leq \frac{1}{t' - (t - r)} (d(\gamma(t - r; x, t), x) + d(x, x')) \leq 2 \left(\frac{1}{r} \int_{t-r}^{t'} \|\dot{\gamma}(s; x, t)\|ds + 1 \right),$$

we obtain that

$$\int_0^{t'} L(\gamma(s; x', t'), v(\gamma(s; x', t'), s), \dot{\gamma}(s; x', t')) ds$$

has a bound depending only on (x, t) and r . By (SL), there exists a constant $M(x, t, r) > 0$ such that

$$\int_0^{t'} \|\dot{\gamma}(s; x', t')\| ds \leq M(x, t, r),$$

where $t' \geq t - r/2 > 0$. It means $\|\dot{\gamma}(s; x', t')\|$ are equi-integrable. Therefore, for (x', t') sufficiently close to (x, t) , there exists a constant $R(x, t, r) > 0$ and $s_0 \in [0, t']$ such that $\|\dot{\gamma}(s_0; x', t')\| \leq R(x, t, r)$. By Lemma A.8, there exists an absolutely continuous function $p(t; x', t')$ satisfying $|p'(t; x', t')| \leq \lambda \|\partial_t v(x, t)\|_\infty$ such that

$$\begin{aligned} & L(\gamma(s; x', t'), v(\gamma(s; x', t'), s), \frac{\dot{\gamma}(s; x', t')}{\theta}) \\ & - L(\gamma(s; x', t'), v(\gamma(s; x', t'), s), \dot{\gamma}(s; x', t')) \geq p(s; x', t')(\theta - 1), \quad \forall \theta > 0. \end{aligned}$$

One can take $\theta = 2$ and $t = s_0$ to obtain the upper bound of $p(s_0)$, and take $\theta = 1/2$ and $t = s_0$ to obtain the lower bound of $p(s_0)$. Note that $p'(t)$ is bounded. We finally obtain the bound of $\|p(t)\|_\infty$, which is independent of (x', t') . Since $L(x, u, \dot{x})$ satisfies (SL), according to Lemma A.9 and taking $\rho = 1$, we have

$$L(\gamma(s; x', t'), v(\gamma(s; x', t'), s), \frac{\dot{\gamma}(s; x', t')}{\|\dot{\gamma}(s; x', t')\|}) \geq \frac{\Theta(\|\dot{\gamma}(s; x', t')\|)}{\|\dot{\gamma}(s; x', t')\|} - \|p(s; x', t')\|_\infty.$$

Therefore, for (x', t') sufficiently close to (x, t) , the minimizers $\gamma(s; x', t')$ have a Lipschitz constant independent of (x', t') .

In order to prove Item (2), we first show that $u(x, t)$ is locally Lipschitz in x . For any $\delta > 0$, fix $(x_0, t) \in M \times [\delta, T]$ and $x, x' \in B(x_0, \delta/2)$. We denote by $d_0 = d(x, x') \leq \delta$ the Riemannian distance between x and x' . Then

$$\begin{aligned} u(x', t) - u(x, t) & \leq \int_{t-d_0}^t L(\alpha(s), v(\alpha(s), s), \dot{\alpha}(s)) ds \\ & \quad - \int_{t-d_0}^t L(\gamma(s; x, t), v(\gamma(s; x, t), s), \dot{\gamma}(s; x, t)) ds, \end{aligned}$$

where $\gamma(s; x, t)$ is a minimizer of $u(x, t)$ and $\alpha : [t - d_0, t] \rightarrow M$ is a geodesic connecting $\gamma(t - d_0; x, t)$ and x' with constant speed. By Lemma 2.3 (1), if $x \in B(x_0, \delta/2)$, the bound of $\|\dot{\gamma}(s; x, t)\|$ depends only on x_0 and δ . Noticing that

$$\|\dot{\alpha}(s)\| \leq \frac{d(\gamma(t - d_0; x, t), x')}{d_0} \leq \frac{d(\gamma(t - d_0; x, t), x)}{d_0} + 1,$$

and

$$d(\gamma(t - d_0; x, t), x) \leq \int_{t-d_0}^t \|\dot{\gamma}(s; x, t)\| ds,$$

the bound of $\|\dot{\alpha}(s)\|$ depends only on x_0 and δ . Exchanging the role of (x, t) and (x', t) , one obtain that

$$|u(x, t) - u(x', t)| \leq J_1 d(x, x'),$$

where J_1 depends only on x_0 and δ . Since M is compact, we conclude that for $t \in (0, T]$, the value function $u(\cdot, t)$ is Lipschitz on M .

We are now going to show the locally Lipschitz continuity of $u(x, t)$ in t . Given $t_0 \geq 3\delta/2$ and $t, t' \in [t_0 - \delta/2, t_0 + \delta/2]$, without any loss of generality, we assume $t' > t$. Then

$$\begin{aligned} u(x, t') - u(x, t) &\leq u(\gamma(t; x, t'), t) - u(x, t) \\ &\quad + \int_t^{t'} L(\gamma(s; x, t'), v(\gamma(s; x, t'), s), \dot{\gamma}(s; x, t')) ds, \end{aligned}$$

where the bound of $\|\dot{\gamma}(s; x, t')\|$ depends only on t_0 and δ . We have shown that for $t \geq \delta$, the following holds

$$u(\gamma(t; x, t'), t) - u(x, t) \leq J_1 d(\gamma(t; x, t'), x) \leq J_1 \int_t^{t'} \|\dot{\gamma}(s; x, t')\| ds \leq J_2(t' - t).$$

Thus, $u(x, t') - u(x, t) \leq J_3(t' - t)$, where J_3 depends only on t_0 and δ . The condition $t' < t$ is similar. We conclude the locally Lipschitz continuity of $u(x, \cdot)$ on $(0, T]$.

At last, we prove Item (3). We first prove that $u(x, t)$ is continuous at $t = 0$. For each $\varphi \in C(M)$, there is a sequence $\varphi_m \in \text{Lip}(M)$ converging to φ uniformly. We take φ and φ_m as the initial functions in (2.7), and denote by $u(x, t)$ and $u_m(x, t)$ the corresponding value functions respectively. Since $v(x, t)$ is fixed, by the non-expansiveness of the Lax-Oleinik semigroup, we have $\|u(x, t) - u_m(x, t)\|_\infty \leq \|\varphi - \varphi_m\|_\infty$. Thus, without loss of generality, we assume the initial function to be Lipschitz in the following discussion. Take a constant curve $\alpha(t) \equiv x$. Let $\gamma : [0, t] \rightarrow M$ be a minimizer of $u(x, t)$. It is obvious that

$$u(x, t) = \varphi(\gamma(0)) + \int_0^t L(\gamma(s), v(\gamma(s), s), \dot{\gamma}(s)) ds \leq \varphi(x) + \int_0^t L(x, v(x, s), 0) ds,$$

so $\limsup_{t \rightarrow 0^+} u(x, t) \leq \varphi(x)$. By (SL), there exists a constant $C > 0$ such that

$$\begin{aligned} \int_0^t L(\gamma(\tau), v(\gamma(\tau), \tau), \dot{\gamma}(\tau)) d\tau &\geq \int_0^t \|\partial_x \varphi\|_\infty \|\dot{\gamma}(\tau)\| d\tau + Ct \\ &\geq \|\partial_x \varphi\|_\infty d(\gamma(0), \gamma(t)) + Ct, \end{aligned}$$

which implies that

$$\int_0^t L(\gamma(\tau), v(\gamma(\tau), \tau), \dot{\gamma}(\tau)) d\tau + \varphi(\gamma(0)) \geq \varphi(x) + Ct.$$

Therefore $\liminf_{t \rightarrow 0^+} u(x, t) \geq \varphi(x)$. Combining with Lemma 2.3 (2), $u(x, t)$ is continuous on $M \times [0, T]$.

By a standard argument, one can show that the value function $u(x, t)$ is a solution of (2.8). We omit the details. □

D Weak KAM solutions and viscosity solutions

Following Fathi [13], one can extend the definitions of backward and forward weak KAM solutions of equation (1.2) by using absolutely continuous calibrated curves instead of C^1 curves.

Definition D.1. A function $u_- \in C(M)$ is called a backward weak KAM solution of (1.2) if the following hold.

(1) For each absolutely continuous curve $\gamma : [t', t] \rightarrow M$, we have

$$u_-(\gamma(t)) - u_-(\gamma(t')) \leq \int_{t'}^t L(\gamma(s), u_-(\gamma(s)), \dot{\gamma}(s)) ds.$$

The above condition reads that u_- is dominated by L and denoted by $u_- \prec L$.

(2) For each $x \in M$, there exists an absolutely continuous curve $\gamma_- : (-\infty, 0] \rightarrow M$ with $\gamma_-(0) = x$ such that

$$u_-(x) - u_-(\gamma_-(t)) = \int_t^0 L(\gamma_-(s), u_-(\gamma_-(s)), \dot{\gamma}_-(s)) ds, \quad \forall t < 0.$$

The curves satisfying the above equality are called $(u_-, L, 0)$ -calibrated curves.

A forward weak KAM solution of (1.2) can be defined in a similar manner. We omit the details.

Proposition D.2. If $u \prec L$, then u is a Lipschitz function on M .

Proof. For each $x, y \in M$, let $\alpha : [0, d(x, y)/\delta] \rightarrow M$ be a geodesic of length $d(x, y)$, with constant speed $\|\dot{\alpha}\| = \delta$ and connecting x and y . Then

$$L(\alpha(s), u(\alpha(s)), \dot{\alpha}(s)) \leq \bar{C} + \lambda \|u\|_\infty, \quad \forall s \in [0, d(x, y)/\delta].$$

Then by $u \prec L$ we have

$$u(y) - u(x) \leq \int_0^{d(x, y)/\delta} L(\alpha(s), u(\alpha(s)), \dot{\alpha}(s)) ds \leq \frac{1}{\delta} (\bar{C} + \lambda \|u\|_\infty) d(x, y).$$

Exchanging the role of x and y , we get the Lipschitz continuity of u . \square

By [13, Corollary 8.3.4], we have

Proposition D.3. Suppose $H(t, x, p)$ is a continuous function, and it is coercive and convex in p , then $u(x, t)$ is a subsolution of $\partial_t u + H(t, x, \partial_x u) = 0$ if $u(x, t)$ is locally Lipschitz and $\partial_t u + H(t, x, \partial_x u) \leq 0$ holds almost everywhere.

Proposition D.4. The following statements are equivalent:

- (1) u_- is a viscosity solution of (E_H) ;
- (2) u_- is a fixed point of T_t^- ;
- (3) u_- is a backward weak KAM solution.

Similarly, the following statements are also equivalent:

- (i) $-u_+$ is a viscosity solution of $H(x, -u, -\partial_x u) = 0$;
- (ii) u_+ is a fixed point of T_t^+ ;
- (iii) u_+ is a forward weak KAM solution.

Proof. By Theorem 1, (2) implies (1). We show that (1) implies (2). Since u_- is a viscosity solution of (E_H) , the function $u(x, t) := u_-(x)$ is the viscosity solution of (CP_H) with the initial condition $u(x, 0) = u_-(x)$. By the comparison theorem, we have $u(x, t) = T_t^- u_-(x)$, which implies $u_- = T_t^- u_-$.

Now we show that (3) implies (2). According to the definition of the backward weak KAM solution, for $u_- \in \mathcal{S}_-$ we have

$$u_-(x) = \inf_{\gamma(t)=x} \left\{ u_-(\gamma(0)) + \int_0^t L(\gamma(\tau), u_-(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau \right\},$$

where the infimum is taken in the class of absolutely continuous curves. We show $u_-(x) \leq T_t^- u_-(x)$. The opposite direction is similar. Assume

$$u_-(x) > T_t^- u_-(x).$$

Let $\gamma : [0, t] \rightarrow M$ with $\gamma(t) = x$ be a minimizer of $T_t^- u_-(x)$. Define

$$F(\tau) := u_-(\gamma(\tau)) - T_\tau^- u_-(\gamma(\tau)).$$

Since $F(t) > 0$ and $F(0) = 0$, there is $s_0 \in [0, t)$ such that $F(s_0) = 0$ and $F(s) > 0$ for $s \in (s_0, t]$. By definition we have

$$T_s^- u_-(\gamma(s)) = T_{s_0}^- u_-(\gamma(s_0)) + \int_{s_0}^s L(\gamma(\tau), T_\tau^- u_-(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau,$$

and

$$u_-(\gamma(s)) \leq u_-(\gamma(s_0)) + \int_{s_0}^s L(\gamma(\tau), u_-(\gamma(\tau)), \dot{\gamma}(\tau)) d\tau,$$

which implies

$$F(s) \leq \lambda \int_{s_0}^s F(\tau) d\tau.$$

By the Gronwall inequality, we conclude $F(s) \equiv 0$ for all $s \in [s_0, t]$, which contradicts $F(t) > 0$.

It remains to show (2) implies (3). For each absolutely continuous curve $\gamma : [t', t] \rightarrow M$, we have

$$\begin{aligned} u_-(\gamma(t)) - u_-(\gamma(t')) &= T_t^- u_-(\gamma(t)) - T_{t'}^- u_-(\gamma(t')) \\ &\leq \int_{t'}^t L(\gamma(s), T_s^- u_-(\gamma(s)), \dot{\gamma}(s)) ds = \int_{t'}^t L(\gamma(s), u_-(\gamma(s)), \dot{\gamma}(s)) ds, \end{aligned}$$

which implies $u_- \prec L$. We now show the existence of the $(u_-, L, 0)$ -calibrated curve. We define a sequence of absolutely continuous curves as follows: Let $\gamma_0(0) = x$ and $\gamma_n : [0, 1] \rightarrow M$ be a minimizer of $T_1^- u_-(\gamma_{n-1}(0))$ with $\gamma_n(1) = \gamma_{n-1}(0)$. We define $\gamma_- : (-\infty, 0] \rightarrow M$ by $\gamma_-(-t) := \gamma_{[t]+1}([t] + 1 - t)$ for all $t > 0$, which is also absolutely continuous. Here, $[t]$ stands for the greatest integer not greater than t . Then we have

$$\begin{aligned} u_-(\gamma_-(-[t])) - u_-(\gamma_-(-t)) &= T_1^- u_-(\gamma_{[t]+1}(1)) - T_{[t]+1-t}^- u_-(\gamma_{[t]+1}([t] + 1 - t)) \\ &= \int_{[t]+1-t}^1 L(\gamma_{[t]+1}(s), T_s^- u_-(\gamma_{[t]+1}(s)), \dot{\gamma}_{[t]+1}(s)) ds \\ &= \int_{-t}^{-[t]} L(\gamma_-(s), u_-(\gamma_-(s)), \dot{\gamma}_-(s)) ds. \end{aligned}$$

Similarly, one can prove that for all $n = 0, 1, \dots$,

$$u_-(\gamma_-(-n)) - u_-(\gamma_-(-n-1)) = \int_{-n-1}^{-n} L(\gamma_-(s), u_-(\gamma_-(s)), \dot{\gamma}_-(s)) ds.$$

We conclude that $\gamma_- : (-\infty, 0] \rightarrow M$ is a $(u_-, L, 0)$ -calibrated curve.

The proof is now complete. \square

References

- [1] Bales, G., An introduction to the theory of viscosity solutions for first-order Hamilton-Jacobi equations and applications, Springer-Heidelberg, 49-109, 2013.
- [2] Ball, J. M. and Mizel, V. J., One-dimensional variational problems whose minimizers do not satisfy the Euler-Lagrange equation, *Arch. Rational Mech. Anal.*, **90**, 1985, 325-388.
- [3] Bettiol, P. and Mariconda, C., A new variational inequality in the Calculus of Variations and Lipschitz regularity of minimizers, *J. Differential Equations*, **5**(268), 2020, 2332-2367.
- [4] Buttazzo, G., Giaquinta, M. and Hildebrandt, S., One-Dimensional Variational Problems, an Introduction, Oxford Lecture Series in Mathematics and its Applications, New York: Clarendon, 1998.
- [5] Cannarsa, P. and Sinestrari, C., Semiconcave functions, Hamilton-Jacobi equations, and optimal control, vol. 58, Springer, New York, 2004.
- [6] Cannarsa, P., Cheng, W., Wang, K. and Yan, J., Herglotz' generalized variational principle and contact type Hamilton-Jacobi equations, *Trends in Control Theory and Partial Differential Equations*, 39-67, Springer INdAM Ser., **32**, Springer, Cham, 2019.
- [7] Cannarsa, P., Cheng, W., Jin, L., Wang, K. and Yan, J., Herglotz' variational principle and Lax-Oleinik evolution, *J. Math. Pures Appl.*, **141**, 2020, 99-136.
- [8] Chen, Q., Cheng, W., Ishii, H. and Zhao, K., Vanishing contact structure problem and convergence of the viscosity solutions, *Comm. Partial Differential Equations*, **44**(9), 2019, 801-836.
- [9] Crandall, M., Ishii, H. and Lions, P.-L., User's guide to viscosity solutions of second order partial differential equations, *Bull. Amer. Math. Soc. (N.S.)*, **27**(1), 1992, 1-67.
- [10] Crandall, M. and Lions, P.-L., Viscosity solutions of Hamilton-Jacobi equations, *Trans. Amer. Math. Soc.*, **277**, 1983, 1-42.
- [11] Crandall, M., Evans, L. and Lions, P.-L., Some properties of viscosity solutions of Hamilton-Jacobi equations, *Trans. Amer. Math. Soc.*, **282**(2), 1984, 487-502.
- [12] Davini, A. and Siconolfi, A., A generalized dynamical approach to the large time behavior of solutions of Hamilton-Jacobi equations, *SIAM J. Math. Anal.*, **38**(2), 2006, 478-502.
- [13] Fathi, A., Weak KAM Theorems in Lagrangian Dynamics. Cambridge: Cambridge University Press, 10th preliminary version, 2008.
- [14] Fathi, A. and Siconolfi, A., PDE aspects of Aubry-Mather theory for quasiconvex Hamiltonians, *Calc. Var. Partial Differential Equations*, **2**, 2005, 185-228.
- [15] Herglotz, G., Berührungstransformationen, Lectures at the University of Göttingen, Göttingen, 1930.
- [16] Hong, J., Cheng, W., Hu, S. and Zhao, K., Representation formulas for contact type Hamilton-Jacobi equations, *J. Dynam. Differential Equations*, Published online.
- [17] Ishii, H., Perron's methods for Hamilton-Jacobi equations, *Duke Math. J.*, **55**, 1987, 369-384.
- [18] Ishii, H., Wang, K., Wang, L. and Yan, J., Hamilton-Jacobi equations with their Hamiltonians depending Lipschitz continuously on the unknown, *Comm. Partial Differential Equations*, **47**, 2022, 417-452.
- [19] Ichihara, N. and Ishii, H., Long-time behavior of solutions of Hamilton-Jacobi equations with convex and coercive Hamiltonians, *Arch. Rational Mech. Anal.*, **194**, 2009, 383-419.
- [20] Jin, L., Wang, L. and Yan, J., A representation formula of viscosity solutions to weakly coupled systems of Hamilton-Jacobi equations with applications to regularizing effect, *J. Differential Equations*, **268**, 2020, 2012-2039.
- [21] Liu, Q., Torres, P. and Wang, C., Contact Hamiltonian dynamics: Variational principles, invariants, completeness and periodic behavior, *Annals of Physics*, **395**, 2018, 26-44.

- [22] Maso, G. D., An introduction to Γ -convergence, Progress in Nonlinear Differential Equations and their Applications, 1993.
- [23] Wang, K., Wang, L. and Yan, J., Implicit variational principle for contact Hamiltonian systems, *Nonlinearity*, **30**, 2017, 492-515.
- [24] Wang, K., Wang, L. and Yan, J., Variational principle for contact Hamiltonian systems and its applications, *J. Math. Pures Appl.*, **123**, 2019, 167-200.
- [25] Wang, K., Wang, L. and Yan, J., Aubry-Mather theory for contact Hamiltonian systems, *Commun. Math. Phys.*, **366**, 2019, 981-1023.
- [26] Wang, K., Wang, L. and Yan, J., Weak KAM solutions of Hamilton-Jacobi equations with decreasing dependence on unknown functions, *J. Differential Equations*, **286**, 2021, 411-432.
- [27] Wang, K., Wang, L. and Yan, J., Aubry-Mather theory for contact Hamiltonian systems II, *Discrete Contin. Dyn. Syst.*, **42**, 2022, 555-595.